

# **GROUND-WATER QUALITY IN GEORGIA FOR 2000**

**John C. Donahue**

GEORGIA DEPARTMENT OF NATURAL RESOURCES  
ENVIRONMENTAL PROTECTION DIVISION  
GEORGIA GEOLOGIC SURVEY

**ATLANTA  
2001**

**CIRCULAR 12P**

# **GROUND-WATER QUALITY IN GEORGIA FOR 2000**

**John C. Donahue**

*The preparation of this report was funded in part through a grant from the U.S. Environmental Protection Agency under the provisions of Section 106 of the Federal Water Pollution Control Act of 1972, as amended.*

GEORGIA DEPARTMENT OF NATURAL RESOURCES  
LONICE C. BARRETT, COMMISSIONER

ENVIRONMENTAL PROTECTION DIVISION  
HAROLD F. REHEIS, DIRECTOR

GEORGIA GEOLOGIC SURVEY  
WILLIAM H. McLEMORE, STATE GEOLOGIST

**ATLANTA  
2001**

**CIRCULAR 12P**

# TABLE OF CONTENTS

<u>Section</u>	<u>Page</u>
<b>CHAPTER 1 <u>INTRODUCTION</u></b>	1-1
1.1 Purpose and Scope	1-1
1.2 Factors Affecting Chemical Ground-Water Quality	1-2
1.3 Hydrogeologic Provinces of Georgia	1-2
1.3.1 Coastal Plain Province	1-4
1.3.2 Piedmont/Blue Ridge Province	1-4
1.3.3 Valley and Ridge Province	1-5
1.4 Regional Ground-Water Problems	1-5
<b>CHAPTER 2 <u>GEORGIA GROUND-WATER MONITORING NETWORK</u></b>	2-1
2.1 Monitoring Stations	2-1
2.2 Uses and Limitations	2-1
2.3 Analyses	2-3
<b>CHAPTER 3 <u>GROUND-WATER QUALITY IN GEORGIA</u></b>	3-1
3.1 Overview	3-1
3.2 Cretaceous Aquifer System	3-1
3.3 Providence Aquifer System	3-4
3.4 Clayton Aquifer System	3-6
3.5 Claiborne Aquifer System	3-9
3.6 Jacksonian Aquifer System	3-12
3.7 Floridan Aquifer System	3-12
3.8 Miocene Aquifer System	3-19
3.9 Piedmont/Blue Ridge Unconfined Aquifers	3-19
3.10 Valley and Ridge Unconfined Aquifers	3-22
<b>CHAPTER 4 <u>SUMMARY AND CONCLUSIONS</u></b>	4-1
<b>CHAPTER 5 <u>LIST OF REFERENCES</u></b>	5-1
<b>APPENDIX</b>	
Laboratory Data	
<b>LIST OF FIGURES</b>	
Figure 1-1 The Hydrogeologic Provinces of Georgia	1-3
Figure 3-1 The Seven Major Aquifer Systems of the Coastal Plain Province	3-2
Figure 3-2 Locations of Stations Monitoring the Cretaceous Aquifer System	3-3
Figure 3-3 Nitrate/Nitrite Concentrations for Selected Wells in the Cretaceous Aquifer System	3-5

<b><u>LIST OF FIGURES (Continued)</u></b>		<b><u>Page</u></b>
Figure 3-4	Locations of Stations Monitoring the Providence Aquifer System	3-7
Figure 3-5	Locations of Stations Monitoring the Clayton Aquifer System	3-8
Figure 3-6	Nitrate/Nitrite Concentrations for Selected Wells in the Clayton Aquifer System	3-10
Figure 3-7	Locations of Stations Monitoring the Claiborne Aquifer System	3-11
Figure 3-8	Nitrate/Nitrite Concentrations for Selected Wells in the Claiborne Aquifer System	3-13
Figure 3-9	Locations of Stations Monitoring the Jacksonian Aquifer System	3-14
Figure 3-10	Nitrate/Nitrite Concentrations for Selected Wells in the Jacksonian Aquifer System	3-15
Figure 3-11	Locations of Stations Monitoring the Floridan Aquifer System	3-17
Figure 3-12	Nitrate/Nitrite Concentrations for Selected Wells in the Floridan Aquifer System	3-18
Figure 3-13	Locations of Stations Monitoring the Miocene Aquifer System	3-20
Figure 3-14	Nitrate/Nitrite Concentrations for Selected Wells in the Miocene Aquifer System	3-21
Figure 3-15	Locations of Stations Monitoring the Piedmont/Blue Ridge Unconfined Aquifers	3-23
Figure 3-16	Nitrate/Nitrite Concentrations for Selected Wells in the Piedmont/Blue Ridge Unconfined Aquifer System: Piedmont Sector	3-24
Figure 3-17	Nitrate/Nitrite Concentrations for Selected Wells in the Piedmont/Blue Ridge Unconfined Aquifer System: Blue Ridge Sector	3-25
Figure 3-18	Locations of Stations Monitoring the Valley and Ridge Unconfined Aquifers	3-26
Figure 3-19	Nitrate/Nitrite Concentrations for Selected Wells and Springs in the Valley and Ridge Unconfined Aquifers	3-28

**LIST OF TABLES**

Table 2-1	Georgia Ground-Water Monitoring Network, Calendar Year 2000	2-2
Table 4-1	Pollution and Contamination Incidents, Calendar Year 2000	4-2
Table A-1	Standard Water Quality Analyses: Anions, Volatile Organic Compounds, and Other Parameters	A-2
Table A-2	Optional Water Quality Analyses: Metals and Anions	A-6
Table A-3	2000 Ground-Water Quality Analyses of the Cretaceous Aquifer System	A-7
Table A-4	2000 Ground-Water Quality Analyses of the Providence Aquifer System	A-10

<b><u>LIST OF TABLES (Continued)</u></b>		<b><u>Page</u></b>
Table A-5	2000 Ground-Water Quality Analyses of the Clayton Aquifer System	A-11
Table A-6	2000 Ground-Water Quality Analyses of the Claiborne Aquifer System	A-12
Table A-7	2000 Ground-Water Quality Analyses of the Jacksonian Aquifer System	A-13
Table A-8	2000 Ground-Water Quality Analyses of the Floridan Aquifer System	A-15
Table A-9	2000 Ground-Water Quality Analyses of the Miocene Aquifer System	A-23
Table A-10	2000 Ground-Water Quality Analyses of the Piedmont/Blue Ridge Aquifer System	A-25
Table A-11	2000 Ground-Water Quality Analyses of the Valley and Ridge Aquifer System	A-29

## **CHAPTER 1 INTRODUCTION**

### **1.1 PURPOSE AND SCOPE**

This report, covering the calendar year 2000, is the sixteenth in a series of summaries discussing the chemical quality of ground water in Georgia. These summaries are among the tools used by the Georgia Environmental Protection Division (EPD) to assess trends in the quality of the State's ground-water resources. EPD is the State organization with regulatory responsibility for maintaining and, where possible, improving ground-water quality and availability. EPD has implemented a comprehensive statewide ground-water management policy of anti-degradation (EPD, 1991; 1998). Five components constitute EPD's ground-water quality assessment program:

1. The Georgia Ground-Water Monitoring Network. The Geologic Survey Branch of EPD maintains this program, which is designed to evaluate the ambient ground-water quality of nine aquifer systems throughout the State of Georgia. The data collected from sampling of the Ground-Water Monitoring Network form the basis for this report.
2. Sampling of public drinking water wells as part of the Safe Drinking Water Program (Water Resources Management Branch). This program provides data on the quality of ground water that the residents of Georgia are using.
3. Special studies addressing specific water quality issues. A survey of nitrite /nitrate levels in shallow wells located throughout the State of Georgia (Shellenberger, et al., 1996; Stuart, et al., 1995) and the operation of a Pesticide Monitoring Network, currently conducted jointly by the Geologic Survey Branch and the Georgia Department of Agriculture (GDA) (Tolford, 1999), are examples of these types of studies.
4. Ground-water sampling at environmental facilities such as municipal solid waste landfills, RCRA facilities, and sludge disposal facilities. The primary agencies responsible for monitoring these facilities are EPD's Land Protection, Water Protection, and Hazardous Waste Management Branches.
5. The development of a wellhead protection program (WHP), which is designed to protect the area surrounding a municipal drinking water well from contaminants. The U.S. Environmental Protection Agency (EPA) approved Georgia's WHP Plan on September 30, 1992. The WHP Plan became a part of the Georgia Safe Drinking Water Rules, effective July 1, 1993. The protection of public water supply wells from contaminants is

important not only for maintaining ground-water quality, but also for ensuring that public water supplies meet health standards.

Analyses of water samples collected for the Georgia Ground-Water Monitoring Network during calendar year 2000 and from previous years form the database for this summary. The Georgia Ground-Water Monitoring Network comprises 128 wells and springs. All stations are generally sampled on an annual basis. Testing for most stations is restricted to volatile organic compounds and nitrate/nitrite. Stations showing recent pollution or contamination may be subject to confirmatory sampling on a basis more frequent than annual. During calendar year 2000, EPD personnel collected 125 samples from 114 wells and 7 springs. A review of the calendar year 2000 data and comparison of these data with those for samples collected as early as 1984 indicate that ground-water quality at most of the 128 sampling sites generally has changed little and remains excellent.

## **1.2 FACTORS AFFECTING CHEMICAL GROUND-WATER QUALITY**

The chemical quality of ground water drawn for sampling is the result of complex physical, chemical, and biological processes. Among the more significant controls are the chemical quality of the water entering the ground-water flow system, the reactions of infiltrating water with the soils and rocks that are encountered, and the effects of the well-and-pump system.

Most water enters the ground-water system in upland recharge areas. Water seeps through interconnected pores and joints in the soils and rocks until discharged to a surface-water body (e.g., stream, river, lake, or ocean). The initial water chemistry, the amount of recharge, and the attenuation capacity of soils have a strong influence on the quality of ground water in recharge areas. Chemical interactions between the water and the aquifer host rocks have an increasing significance with longer underground residence times. As a result, ground water from discharge areas tends to be more highly mineralized than ground water in recharge areas.

The well-and-pump system can also have a strong influence on the quality of the well water. Well casings, through compositional breakdown, can contribute metals (e.g., iron from steel casings) and organic compounds (e.g., tetrahydrofuran from PVC pipe cement) to the water. Pumps often aerate the water being discharged. An improperly constructed well can present a conduit that allows local pollutants to enter the ground-water flow system.

## **1.3 HYDROGEOLOGIC PROVINCES OF GEORGIA**

This report defines three hydrogeologic provinces in Georgia by their general geologic and hydrologic characteristics (Figure 1-1). These provinces consist of:



Figure 1-1 The Hydrogeologic Provinces of Georgia



1. the Coastal Plain Province of south Georgia;
2. the Piedmont/Blue Ridge Province, which includes all but the northwest corner of north Georgia; and
3. the Valley and Ridge Province of northwest Georgia.

### **1.3.1 Coastal Plain Province**

Georgia's Coastal Plain Province generally comprises a wedge of loosely consolidated sediments that gently dip and thicken to the south and southeast. Ground water in the Coastal Plain Province flows through interconnected pore space between grains in the host rocks and through solution-enlarged voids.

The oldest outcropping sedimentary formations (Cretaceous) are exposed along the Fall Line, which is the northern limit of the Coastal Plain Province. Successively younger formations occur at the surface to the south and southeast.

The Coastal Plain contains Georgia's major confined (artesian) aquifers. Confined aquifers are those in which a layer of impermeable material (i.e., clay or shale) holds the top of the water column below the level to which it would normally rise. Water enters the aquifers in their updip outcrop areas, where the more permeable sediments of the aquifer tend to be exposed. Many Coastal Plain aquifers are unconfined in their updip outcrop areas, but become confined in downdip areas to the southeast, where they are overlain by successively younger rock formations. Ground-water flow through confined Coastal Plain aquifers is generally to the south and southeast, in the direction of the dip of the rocks.

The sediments forming the seven major aquifers in the Coastal Plain range in age from Cretaceous to Miocene. Horizontal and vertical changes in the permeability of the rock units that form these aquifers determine the thickness and extent of the aquifers. Several aquifers may be present in a single geographic area, forming a vertical "stack".

The Cretaceous and Jacksonian aquifer systems (primarily sands) are a common source of drinking water within a 35-mile wide band that lies adjacent to and south of the Fall Line. Southwestern Georgia relies on four vertically stacked aquifers (sands and carbonates) for drinking-water supplies: the Providence, Clayton, Claiborne and Floridan aquifer systems. The Floridan aquifer system (primarily carbonates) serves most of south-central and southeastern Georgia. The Miocene aquifer system (primarily sands) is the principal "shallow" unconfined aquifer system occupying much of the same broad area underlain by the Floridan aquifer system. It becomes confined in the coastal counties and locally in the Grady, Thomas, Brooks and Lowndes County area of south Georgia.

### **1.3.2 Piedmont/Blue Ridge Province**

Crystalline rocks of metamorphic and igneous origin (primarily Precambrian and Paleozoic in age) underlie the Piedmont and Blue Ridge Provinces. These two provinces differ geologically but are discussed together here because they share common hydrologic

properties. The principal water-bearing features are fractures, compositional layers, and other geologic discontinuities in the rock, as well as intergranular porosity in the overlying soil and saprolite horizons. Thick soils and saprolites are often important as the “reservoir” that supplies water to the water-bearing fracture and joint systems. Ground water typically flows from local highlands toward discharge areas along streams. However, during prolonged dry periods or in areas of heavy pumpage, surface water may flow from the streams into the ground-water systems.

### **1.3.3 Valley and Ridge Province**

Consolidated Paleozoic sedimentary formations characterize the Valley and Ridge Province. The principal permeable features of the Valley and Ridge Province are fractures and solution voids; intergranular porosity also is important in some places. Locally, ground-water and surface-water systems closely interconnect. Dolostones and limestones of the Knox Group are the principal aquifers where they occur in the axes of broad valleys. The greater hydraulic conductivities of the thick carbonate sections in this Province, in part due to solution-enlarged joints, permit development of higher yielding wells than in the Piedmont and Blue Ridge Province.

## **1.4 REGIONAL GROUND-WATER PROBLEMS**

Data from ground-water investigations in Georgia, including those from the Ground-Water Monitoring Network, indicate that virtually all of Georgia has shallow ground water sufficient for domestic supply. Iron, aluminum, and manganese are the only constituents that occur routinely in concentrations exceeding drinking-water standards. These metals are naturally occurring and do not pose a health risk. Iron and manganese can cause reddish to brownish stains on objects.

Only a few occurrences of polluted or contaminated ground waters are known from North Georgia (see Section 4). Aquifers in the outcrop areas of Cretaceous sediments south of the Fall Line typically yield acidic water that may require treatment. The acidity occurs naturally and results both from the inability of the sandy aquifer sediments to neutralize acidic rainwater and from biologically influenced acid-producing reactions between infiltrating water and soils and sediments. Nitrite/nitrate concentrations in shallow ground water from the farm belt of southern Georgia are usually within drinking-water standards, but are somewhat higher than levels found in other areas of the State.

Besides the karst plain area (Dougherty Plain) in southwest Georgia, the Floridan aquifer system contains two other areas of naturally-occurring reduced ground-water quality. The first is the area of the Gulf Trough, a narrow, linear geological feature extending from southwestern Deatur County through central Bulloch County. Here, ground water is typically high in total dissolved solids and contains elevated levels of barium, sulfate, and radionuclides. The second is the coastal area of Georgia, where influx of

water with high dissolved solids contents presents problems. In the Brunswick area, ground-water withdrawal from the upper Floridan results in up-coning of water with high dissolved solids contents from deeper parts of the aquifer. In the Savannah region, a cone of depression caused by pumping in and around Savannah induces saline ground water to flow down-gradient from the Port Royal Sound area of South Carolina toward Savannah.

## **CHAPTER 2 GEORGIA GROUND-WATER MONITORING NETWORK**

### **2.1 MONITORING STATIONS**

Stations of the calendar year 2000 Ground-Water Monitoring Network are situated in the seven major aquifers and aquifer systems of the Coastal Plain Province and in the unconfined ground-water systems of the Piedmont and Blue Ridge Provinces and of the Valley and Ridge Province (Table 2-1). Monitoring stations are located in three critical settings:

1. areas of surface recharge;
2. areas of potential pollution related to regional activities (e.g., agricultural and industrial areas); and
3. areas of significant ground-water use.

Most of the monitoring stations are municipal, industrial, and domestic wells that have reliable well-construction data. The Monitoring Network also includes monitoring wells in specific areas where the State's aquifers are recognized to be especially susceptible to contamination or pollution (e.g., the Dougherty Plain of southwestern Georgia and the State's coastal area).

### **2.2 USES AND LIMITATIONS**

Regular sampling of wells and springs of the Ground-Water Monitoring Network permits analysis of ground-water quality with respect to location (spatial trends) and with respect to the time of sample collection (temporal trends). Spatial trends are useful for assessing the effects of the geologic framework of the aquifer and regional land-use activities on ground-water quality. Temporal trends permit an assessment of the effects of rainfall and drought periods on ground-water quantity and quality. Both trends are useful for the detection of non-point source pollution. Non-point source pollution arises from broad-scale phenomena such as acid rain deposition and application of agricultural chemicals on crop lands.

It should be noted that the data of the Ground-Water Monitoring Network represent water quality in only limited areas of Georgia. Monitoring water quality at 128 sites located throughout Georgia provides an indication of ground-water quality at the locality sampled and at the horizon corresponding to the screened interval in the well or to the head of the spring at each station in the Monitoring Network. Caution should be exercised in drawing strict conclusions and applying any results reported in this study to ground waters that are not being monitored.

Stations of the Ground-Water Monitoring Network intentionally are located away from known point sources of pollution. The wells provide baseline data on ambient water

Table 2-1. Georgia Ground-Water Monitoring Network, Calendar Year 2000.

AQUIFER SYSTEM	NUMBER OF MONITORING STATIONS VISITED & SAMPLES TAKEN, YEAR 2000	PRIMARY STRATIGRAPHIC EQUIVALENTS	AGE OF AQUIFER FORMATIONS
Cretaceous	17 stations (18 samples)	Ripley Formation, Cusseta Sand, Blufftown Formation, Eutaw Formation, Tuscaloosa Formation, Steel Creek Formation, Gaillard Formation, Pio Nono Formation	Late Cretaceous
Providence	2 stations (2 samples)	Providence Sand	Late Cretaceous
Clayton	5 stations (5 samples)	Clayton Formation	Paleocene
Claiborne	5 stations (5 samples)	Claiborne Group	Middle Eocene
Jacksonian	8 stations (8 samples)	Barnwell Group	Late Eocene
Floridan	48 stations (48 samples)	Predominantly Suwannee Limestone and Ocala Group	Predominantly Middle Eocene to Oligocene
Miocene	7 stations (7 samples)	Predominantly Altamaha Formation and Hawthorne Group	Miocene-Recent
Piedmont/Blue Ridge	20 stations (23 samples)	Various igneous and metamorphic complexes	Predominately Paleozoic and Precambrian
Valley and Ridge	9 stations (9 samples)	Shady Dolomite, Knox Group, and Chickamauga Group	Paleozoic: Cambrian and Ordovician

quality in Georgia. EPD requires other forms of ground-water monitoring for activities that may result in point source pollution (e.g., landfills, hazardous waste facilities and land application sites) through its environmental facilities permit programs.

Ground-water quality changes gradually and predictably in the areally extensive aquifers of the Coastal Plain Province. The Monitoring Network allows for some definition of the chemical processes occurring in large confined aquifers. Unconfined aquifers in northern Georgia and the surface recharge areas of southern Georgia are of comparatively small areal extent and more open to interactions with land-use activities. The wide spacing of monitoring stations does not permit equal characterization of water-quality processes in these settings. The quality of water from monitoring wells completed in unconfined aquifers represents only the general nature of ground water in the vicinity of the monitoring wells. Ground water in the recharge areas of the Coastal Plain aquifers is the future drinking-water resource for down-flow areas. Monitoring wells in these recharge areas, in effect, constitute an early warning system for potential future water quality problems in confined portions of the Coastal Plain aquifers.

### **2.3 ANALYSES**

Analyses are available for 125 water samples collected during calendar year 2000 from 121 stations (114 wells and 7 springs). In 1984, the first year of the Ground-Water Monitoring Network, hydrogeologists sampled water from 39 wells in the Piedmont/Blue Ridge and Coastal Plain Provinces. Since 1984, the Ground-Water Monitoring Network has been expanded through addition of further wells and springs to cover all three hydrogeologic provinces, with most of the monitoring performed in the Coastal Plain.

Ground water from all monitoring stations is tested for nitrate/nitrite and volatile organic compounds (VOCs) including methyl-tert-butyl ether (MTBE). Testing for metals and select anions that are subject to Primary Maximum Contaminant Levels (MCLs) continues for stations that have shown past contamination by these substances. Before collecting a sample, EPD personnel also observe and record certain field parameters -- pH, electrical conductivity, dissolved oxygen, and temperature -- using field instruments. This Circular reports the pH, conductivity, and the chemical analysis results. Files at the Geologic Survey Branch contain the records of all the field parameter measurements and chemical analyses.

The Drinking Water Program of the EPD's Water Resources Management Branch has established MCLs for certain parameters included in the analyses done on Ground-Water Monitoring Network samples (EPD, 1997). Primary MCLs pertain to parameters that may have adverse effects on human health when their values are exceeded. Secondary MCLs pertain to parameters that may give drinking water objectionable, though not health-threatening, properties that may cause persons served by public water systems to cease its use. Foul odor and unpleasant taste are examples of such properties. MCLs apply only to treated water offered for public consumption; nevertheless, they are

useful guidelines for evaluating the quality of untreated (raw) water. Tables A-1 and A-2 in the Appendix list the Primary and Secondary MCLs for Ground Water Monitoring Network parameters.

Most of the wells originally on the Monitoring Network had in-place pumps. Using such pumps to purge the wells and collect samples reduces the potential for cross-contamination of wells. For those wells that lacked in-place pumps, EPD personnel used portable pumps for purging and sampling. In recent years, however, all wells that lacked in-place pumps were dropped from the Monitoring Network, except for a flowing well tapping the lower Floridan, GWN-PA9C.

Sampling procedures are adapted from techniques used by the USGS and the EPA. Hydrogeologists purge the wells (three to five times the volume of the water column in the well) before the collection of a sample to reduce the influence of the well, pump and distribution system on water quality. Municipal, industrial, and domestic wells typically require approximately 30 to 45 minutes of purging before sample collection.

During purging, a manifold captures flow at the pump system discharge point before the water is exposed to the atmosphere and conducts it past the field instrument probes, while EPD personnel observe (and record) the field parameters. With increased purging time, typical trends for field parameters include a lowering of pH, dissolved oxygen content, and conductivity, and a transition toward the mean annual air temperature. The hydraulic flow characteristics of unconfined aquifers, the depth of withdrawal, and pump effects may alter these trends.

Samples are collected once the field parameters stabilize or otherwise indicate that the effects of the well have been minimized. EPD personnel fill the sample bottles and then promptly place them on ice to preserve the water quality. The personnel next transport the samples to the laboratories for analysis on or before the Friday of the week in which they were collected.

## **CHAPTER 3 GROUND-WATER QUALITY IN GEORGIA**

### **3.1 OVERVIEW**

Georgia's nine major aquifers and aquifer systems are grouped into three hydrogeologic provinces for the purposes of this report.

The Coastal Plain Province comprises seven major aquifers or aquifer systems that are restricted to specific regions and depths within the province (Figure 3-1). These major aquifer systems commonly incorporate smaller aquifers that are locally confined. Ground-Water Monitoring Network wells in the Coastal Plain aquifers are generally located in three settings:

1. Recharge (or outcrop) areas that are located in regions that are geologically updip and generally to the north of confined portions of these aquifers.
2. Updip, confined areas that are located in regions that are proximal to the recharge areas, yet are confined by overlying geologic formations. These areas are generally south to southeast of the recharge areas.
3. Downdip, confined areas, located to the south and southeast in the deeper, confined portions of the aquifers distal to the recharge areas.

Small-scale, localized ground-water flow patterns characterize the two hydrogeologic provinces of north Georgia, the Piedmont/Blue Ridge Province and the Valley and Ridge Province. Deep regional flow systems are unknown in northern Georgia. Geologic discontinuities (such as fractures) and compositional changes within the aquifer generally control ground-water flow in the Piedmont/Blue Ridge Province. Local topographic features, such as hills and valleys, influence ground-water flow patterns. Many of the factors controlling ground-water flow in the Piedmont/Blue Ridge Province also apply in the Valley and Ridge Province. The Valley and Ridge Province additionally possesses widespread karst features, which significantly enhance porosity and permeability in localized areas and exert a strong influence on local ground-water flow patterns.

### **3.2 CRETACEOUS AQUIFER SYSTEM**

The Cretaceous aquifer system is a complexly interconnected group of aquifer subsystems developed in the Late Cretaceous sands of the Coastal Plain Province. These sands crop out in an extensive recharge area immediately south of the Fall Line in west and central Georgia (Figure 3-2). Overlying Tertiary sediments restrict Cretaceous outcrops to valley bottoms in parts of the northeastern Coastal Plain. Five distinct subsystems of the Cretaceous aquifer system, including the Providence aquifer system, are recognized west of the Ocmulgee River (Pollard and Vorhis, 1980). These merge into






Figure 3-1 The Seven Major Aquifer Systems of the Coastal Plain Province




Figure 3-2 Locations of Stations Monitoring the Cretaceous Aquifer System

three subsystems to the east (Clarke, et al., 1985; Huddlestun and Summerour, 1996). Aquifer sands thicken south-ward from the Fall Line, from where they pinch out against crystalline Piedmont rocks, to a sequence of sand and clay approximately 2,000 feet thick at the southern limits of the main aquifer-use area (limit of utilization, Figure 3-2). Vertical leakage from overlying members of the aquifer system provides significant recharge in downdip areas.

EPD collected 18 samples from 17 wells in calendar year 2000 to monitor the water quality of the Cretaceous aquifer system, exclusive of the Providence aquifer system (Figure 3-2). Table A-3 in the Appendix lists the analytical results for samples collected from the Cretaceous aquifer system. Four of the sampled wells, GWN-K8, GWN-K9, GWN-K12, and GWN-K20 are located away from the Cretaceous outcrop and recharge area, while the remainder lie within the general recharge area. The pHs of the sampled waters ranged from 4.21 to 9.22, with the majority (14) being acidic. Electrical conductivities ranged from 13 to 372 microsiemens (uS)/cm, with the lowest occurring in waters from recharge area wells.


Water samples from all 17 wells received testing for nitrite/nitrate and for volatile organic compounds (VOCs). Twelve wells yielded samples with detectable nitrate/ nitrite, with the highest concentration, 0.95 ppm as nitrogen, occurring in a sample from well GWN-K5. Figure 3-3 shows trends in levels of nitrate/nitrite (reported as parts per million [ppm] nitrogen) for selected wells.

Three wells, all located in industrial settings, yielded samples containing VOCs. Well GWN-K1 yielded a sample containing trichloroethylene (0.64 ppb) and chloroform (1.5 ppb). Well GWN-K5 gave a sample containing trichloroethylene (0.88 ppb). As the well is used as a public supply well and had, during the previous year, yielded a sample containing the compound, a follow-up sampling was undertaken. The follow-up sample also contained trichloroethylene (1.1 ppb). A sample from well GWN-K16 contained the trihalomethanes chloroform (4.2 ppb) and bromodichloromethane (1.5 ppb). All VOC concentrations are below primary MCLs.

For well GWN-K1, regular testing for VOCs did not begin until 1999. Before that year, VOC testing had been performed twice, with one occasion finding low-level pollution by trichloroethylene and 1,2-dichloroethylene and the other finding none. Well GWN-K5 has been tested regularly for VOCs since 1993, but has experienced pollution by VOCs only during 1999 and 2000. Well GWN-K16 had been overhauled and had just been returned to service the day before the sampling date. Disinfectants used during the overhaul may have introduced the trihalomethanes found in the sample from that well. The well has no prior history of VOC contamination.

### **3.3 PROVIDENCE AQUIFER SYSTEM**

Sand and coquinoid limestones of the Late Cretaceous Providence Formation comprise the Providence aquifer system of southwestern Georgia. Outcrops of the aquifer



Nitrate/nitrite levels below the detection limit are assigned a value of 0.01 ppm. A missing bar indicates that samples were not collected for that year.

Figure 3-3 Nitrate/Nitrite Concentrations for Selected Wells in the Cretaceous Aquifer System

system extend from northern Clay and Quitman Counties through eastern Houston County (Figure 3-4). At its updip extent, the aquifer system thickens both to the east and to the west of a broad area adjacent to the Flint River. The aquifer system also generally thickens downdip, with an area where the thickness exceeds 300 feet existing in Pulaski County and an area of similar thickness indicated in the Baker/Calhoun/Early county region (Clarke, et al., 1983). Figure 3-4 also shows the downdip limit of the area in which the aquifer system is utilized.

The permeable Providence Formation-Clayton Formation interval forms a single aquifer in the updip areas (Long, 1989) and to the east of the Flint River (Clarke, et al., 1983). This same interval is recognized as the Dublin aquifer system to the east of the Ocmulgee River (Clarke, et al., 1985). Outcrop areas and adjacent covered areas to the east of the Flint River, where the aquifer is overlain by permeable sand units, are surface recharge areas. The Chattahoochee River forms the western discharge boundary for this flow system in Georgia.

EPD sampled two wells drawing from the Providence aquifer system in calendar year 2000. Well GWN-PD2B, a recharge area well, yielded acidic water with a low electrical conductivity and a detectable nitrate/nitrite content. Well GWN-PD3, a downdip well, yielded basic water with a moderate electrical conductivity and no detectable nitrate/nitrite. No VOCs were found in the samples from either well. Analysis results are given in Table A-4 in the Appendix.

### **3.4 CLAYTON AQUIFER SYSTEM**

The Clayton aquifer system of southwestern Georgia is developed mainly in the middle limestone unit of the Paleocene Clayton Formation. Limestones and calcareous sands of the Clayton aquifer system crop out in a narrow belt extending from northeastern Clay County to southwestern Schley County (Figure 3-5). Aquifer thickness varies, ranging from 50 feet near outcrop areas to 265 feet in southeastern Mitchell County (Clarke, et al., 1984). Both the Flint River, to the east, and the Chattahoochee River, to the west, are areas of discharge for the aquifer system in its updip extent. Leakage from the underlying Providence aquifer system and from permeable units in the overlying Wilcox confining zone provides significant recharge in downdip areas (Clarke, et al., 1984). The Clayton and Providence Formations merge to form a single aquifer unit in updip areas (Long, 1989) as well as east of the Flint River (Clarke, et al., 1983). West of the Flint River and downdip, the Clayton/Providence confining zone, a silt and clay-bearing interval, confines the aquifer below (McFadden and Perriello, 1983). In the area east of the Ocmulgee River, the combination of these two aquifers is referred to as the Dublin aquifer system (Clarke, et al., 1985). Figure 3-5 also shows the downdip limit of the area in which the aquifer system is used.

During calendar year 2000, EPD collected five water samples from five wells to monitor the water quality in the Clayton aquifer system (Figure 3-5). Three wells (GWN-




Figure 3-4 Locations of Stations Monitoring the Providence Aquifer System




Figure 3-5 Locations of Stations Monitoring the Clayton Aquifer System

CT5A, GWN-CT7A, GWN-CT8) are located in or near the recharge area, with the latter two wells being less than 100 feet deep. Wells GWN-CT2A and GWN-CT3 were used to sample the downdip portion of the aquifer system.

The pH of the waters from the Clayton wells ranged from acidic to slightly basic. The samples were analyzed for VOCs and nitrate/nitrite. The two shallow recharge area wells yielded waters with the lower electrical conductivities. Nitrate/nitrite levels ranged from undetected to 3.6 ppm as nitrogen. The sample with the highest nitrate/nitrite level came from well GWN-CT7A, a shallow updip well located near a livestock enclosure. The enclosure was unoccupied, a factor possibly contributing to a lower nitrate/nitrite level than in the past. Figure 3-6 shows trends in nitrate/nitrite concentrations for selected wells in the Clayton aquifer system. No VOCs were detected in any of the samples. Table A-5 in the Appendix lists analyses for water samples from the Clayton wells.

### **3.5 CLAIBORNE AQUIFER SYSTEM**


The Claiborne aquifer system is developed primarily in the sandy units in the middle and lower portions of the Middle Eocene Claiborne Group of southwestern Georgia. Claiborne Group sands crop out in a belt extending from northern Early County through western Dooly County. Recharge to the aquifer system occurs both as direct infiltration of precipitation in the recharge area and as leakage from the overlying Floridan aquifer system (Hicks, et al., 1981; Gorday, et al., 1997). Discharge boundaries of the aquifer system are the Ocmulgee River to the east, and the Chattahoochee River to the west. The aquifer is more than 350 feet thick near its downdip limit of utilization (Figure 3-7) (Tuohy, 1984).

The aquifer generally thickens from the outcrop area toward the southeast. The clay-rich upper portion of the Claiborne Group, the Lisbon Formation, acts as a confining layer and separates the aquifer from the overlying Floridan aquifer (McFadden and Perriello, 1983; Long, 1989; Huddleston and Summerour, 1996). The lower water-bearing parts of the group had been correlated to the Tallahatta Formation (e.g., McFadden and Perriello, 1983; Long, 1989; Clarke et al., 1996) or, more recently, have been divided into two formations, the upper one termed the Still Branch Sand and the lower one correlated to the Congaree Formation (Huddleston and Summerour, 1996). The permeable lower units are included in the Gordon aquifer system east of the Ocmulgee River (Brooks, et al., 1985).

During calendar year 2000, EPD personnel drew five samples from five wells to monitor the water quality of the Claiborne aquifer system. Wells GWN-CL2, GWN-CL4A, and GWN-CL8 lie within or near the recharge area, and wells GWN-CL6 and GWN-CL9 tap the downdip portion of the aquifer system, near the limit of utilization.

Two of the recharge area wells yielded acidic water, while one recharge area well and the two downdip wells yielded basic water. The lowest conductivity was measured





Nitrate/nitrite levels below the detection limit are assigned a value of 0.01 ppm. A missing bar indicates that samples were not collected for that year.

Figure 3-6 Nitrate/Nitrite Concentrations for Selected Wells in the Clayton Aquifer System




Figure 3-7 Locations of Stations Monitoring the Claiborne Aquifer System

at an updip well (GWN-CL4A), while the highest was measured in a downdip well (GWN-CL6). All samples were analyzed for VOCs and none were detected. All samples were also analyzed for nitrate/nitrite, which was detected in two of the recharge area samples. Figure 3-8 shows trends in nitrate/nitrite concentrations for selected wells, and Table A-6 in the Appendix gives the analytical results for the Claiborne wells.

### **3.6 JACKSONIAN AQUIFER SYSTEM**


The Jacksonian aquifer system of central and east-central Georgia is predominantly comprised of sands of the Eocene Barnwell Group, though locally, isolated limestone bodies are important. Barnwell Group outcrops extend from Macon and Crawford Counties (Hetrick, 1990) eastward to Burke and Richmond Counties (Hetrick, 1992). Figure 3-9 shows the most significant Jacksonian recharge areas. Aquifer sands form a northern clastic facies of the Barnwell Group; the sands grade southward into less permeable silts and clays of a transition facies (Vincent, 1982). The water-bearing sands are relatively thin, ranging from ten to fifty feet in thickness. Limestones equivalent to the Barnwell Group form a southern carbonate facies and are included in the Floridan aquifer system. The Savannah River and Ocmulgee River are eastern and western discharge boundaries respectively for the updip flow system of the Jacksonian aquifer system. The Jacksonian aquifer system is equivalent to the Upper Three Runs aquifer as used in Summerour et al. (1994).

EPD monitored the water quality of the Jacksonian aquifer system in calendar year 2000 by taking eight samples from eight wells (Figure 3-9). Six wells are in the clastic facies (one, GWN-J2B, drawing from an isolated limestone body), and two wells (GWN-J3 and GWN-J5) are in the transition facies. The pH of the sampled water ranged from 4.61 to 7.56. Conductivity measurements were lowest for two shallow updip clastic facies wells GWN-J7 and GWN-J8. Table A-7 in the Appendix lists the analytical results for all the Jacksonian aquifer wells sampled.

All samples received tests for nitrate/nitrite and for VOCs. No VOCs were detected. Nitrate/nitrite concentrations ranged from undetectable to 8.5 ppm as N. The considerably elevated nitrate/nitrite level of 8.5 ppm as N occurred in the sample from well GWN-J8, a shallow domestic well, which is located near a row crop field. The well has a history of yielding high nitrate/nitrite water samples. Figure 3-10 depicts trends in nitrite/nitrate concentrations for selected wells. The sample from well GWN-J8 was analyzed for beryllium, because of occasional past findings, and 4.3 ppb beryllium was detected (above the primary MCL of 4 ppb). The source of the beryllium remains unknown and is presumed natural.

### **3.7 FLORIDAN AQUIFER SYSTEM**

The Floridan aquifer system consists predominantly of Eocene and Oligocene limestones and dolostones that underlie most of the Coastal Plain Province. The aquifer is



Nitrate/nitrite levels below the detection limit are assigned a value of 0.01 ppm. A missing bar indicates that samples were not collected for that year.

Figure 3-8 Nitrate/Nitrite Concentrations for Selected Wells in the Claiborne Aquifer System





Figure 3-9 Locations of Stations Monitoring the Jacksonian Aquifer System



A missing bar indicates that samples were not collected for that year.

Figure 3-10 Nitrate/Nitrite Concentrations for Selected Wells in the Jacksonian Aquifer System

a major source of ground water for much of its outcrop area and throughout its downdip extent to the south and east.

The upper water-bearing units of the Floridan are the Eocene Ocala Group and the Oligocene Suwanee Limestone (Crews and Huddleston, 1984). These limestones crop out in the Dougherty Plain (a karstic area in southwestern Georgia) and in adjacent areas along a strike to the northeast. In Camden and Wayne counties the Oligocene unit is absent, and the upper part of the Floridan is restricted to units of Eocene age (Clarke, et al., 1990). The lower portion of the Floridan consists mainly of dolomitic limestone of middle and early Eocene age and pelletal, vuggy, dolomitic limestone of Paleocene age, but extends into the late Cretaceous in Glynn County. The lower Floridan is deeply buried and not widely used, except in several municipal and industrial wells in the Savannah area (Clarke, et al., 1990). From its updip limit, defined in the east by clays of the Barnwell Group, the aquifer thickens to well over 700 feet in coastal Georgia. A dense limestone facies along the trend of the Gulf Trough locally limits ground-water quality and availability (Kellam and Gorday, 1990). The Gulf Trough is a linear depositional feature in the Coastal Plain that extends from southwestern Decatur County through central Bulloch County.

A ground-water divide separates a smaller southwestward flow regime in the Floridan aquifer system in the Dougherty Plain from the larger southeastward flow regime in the remainder of Georgia. Rainfall infiltration in outcrop areas and downward leakage from extensive surficial residuum recharge the Dougherty Plain flow system (Hayes, et al., 1983). The main body of the Floridan aquifer system, to the east, is recharged by leakage from the Jacksonian aquifer system and by rainfall infiltration in outcrop areas and in areas where overlying strata are thin. Significant recharge also occurs in the area of Brooks, Echols and Lowndes counties, where the Withlacoochee River and numerous sinkholes breach upper confining beds (Krause, 1979).

During calendar year 2000, EPD collected 48 samples from 48 wells in the Floridan aquifer system (Figure 3-11). All samples underwent testing for nitrate/nitrite and for VOCs. The pH values for all sampled waters were basic. Conductivities ranged from 143 uS/cm to 1279 uS/cm. Most of the wells yielding water with higher conductivity are deeper ones located along the coast. Table A-8 in the Appendix lists the analytical results for the Floridan wells.

Two wells yielded samples with low-level pollution by VOCs: GWN-PA17 and GWN-PA33A. The offending substances in each case were trihalomethanes, which may result from the reflux of treated (chlorinated) water back into the well bore. In each case, the concentrations were well below the primary MCL for trihalomethanes (100 ppb).

Nitrate/nitrite concentrations ranged from undetectable to 4.5 ppm as nitrogen. With one exception, the wells yielding water with the highest nitrate/nitrite contents are located in the Dougherty Plain. Figure 3-12 shows trends in nitrate concentrations for selected Floridan wells.





Figure 3-11 Locations of Stations Monitoring the Floridan Aquifer System





A missing bar indicates that samples were not available for that year.

Figure 3-12 Nitrate/Nitrite Concentrations for Selected Wells in the Floridan Aquifer System

### **3.8 MIOCENE AQUIFER SYSTEM**

Much of south-central and southeastern Georgia lies within outcrop areas of the Miocene Altamaha Formation and Hawthorne Group. Discontinuous lens-shaped bodies of sand, 50 to 80 feet thick, are the main permeable units. Miocene clays and sandy clays are thickest, more than 500 feet, in Wayne County (Watson, 1982).

Areas of confinement exist in the coastal counties. Leakage from overlying surface aquifers into the Miocene aquifer system and, in some areas, from the underlying Floridan aquifer system is significant in the coastal counties (Watson, 1982). Here, two principal aquifer units are present (Joiner, et al., 1988). Clarke (et. al., 1990) use the names upper and lower Brunswick aquifers to refer to these two sandy aquifer units.

EPD collected seven water samples from seven wells to monitor the water quality in the Miocene aquifer system (Figure 3-13). The pH of the samples ranged from 3.80 to 7.28, with six stations producing acidic water. Conductivities ranged from 57 uS/cm to 239 uS/cm. Table A-9 in the Appendix lists the analytical results for Miocene samples.

Nitrate/nitrite data are available for all seven stations. Concentrations ranged from undetected to 27 ppm as nitrogen. Only one well, GWN-MI15, a domestic well, produced a sample with a concentration in excess of the primary MCL of 10 ppm as nitrogen. However, three other wells, GWN-MI5, GWN-MI7, and GWN-MI9A, gave samples with nitrate/nitrite concentrations that were elevated (herein set at the 5 ppm “trigger” level for public water supplies). All three wells are shallow domestic-type wells. Wells GWN-MI7, GWN-MI9A, and GWN-MI15 lie near row crop fields, while well GWN-MI5 is located near a newly installed and occupied animal enclosure. The field near GWN-MI9A, left fallow during 1999, was planted in corn during 2000. Fertilizing the field incident to raising corn may have contributed to the approximate 5 ppm increase of nitrate/nitrite in the sampled water for the year 2000 over that for the year 1999. Figure 3-14 illustrates trends in nitrate/nitrite concentrations for selected wells drawing from the Miocene aquifer system. VOC tests were performed for all seven samples and none contained detectable VOCs.

### **3.9 PIEDMONT/BLUE RIDGE UNCONFINED AQUIFERS**

Georgia's Piedmont and Blue Ridge Physiographic Provinces are developed on metamorphic and igneous rocks that are predominantly Precambrian and Paleozoic in age. Soil and saprolite horizons, compositional layers, and openings along fractures and joints in the rocks are the major water-bearing features. Fracture density and interconnection provide the primary controls on the rate of water flow into wells completed in crystalline rocks. The permeability and thickness of soils and saprolite horizons determine the amount of well yield that can be sustained.

EPD collected twenty-three samples from seventeen wells and three springs to monitor water quality in the Piedmont/Blue Ridge unconfined aquifers. Figure 3-15





Figure 3-13 Locations of Stations Monitoring the Miocene Aquifer System



A missing bar indicates that samples were not collected for that year.

Figure 3-14 Nitrate/Nitrite Concentrations for Selected Wells in the Miocene Aquifer System

shows the locations of the monitoring stations. The pH of the water samples ranged from 4.83 to 7.96, with the majority of the stations yielding slightly acidic water. Conductivities ranged from 18 uS/cm to 359 uS/cm.

All samples were tested for nitrate/nitrite and for VOCs. Because of its history of giving samples with high fluoride concentrations, the sample from station GWN-P12A received testing for inorganic anions besides nitrate/nitrite. An analytical summary for the Piedmont/Blue Ridge sampling stations is in Appendix Table A-10.

Nitrate/nitrite concentrations ranged from undetectable to 1.6 ppm as nitrogen (the primary MCL is 10 ppm as nitrogen). Figures 3-16 and 3-17 show nitrite/nitrate concentrations in selected stations from the Piedmont and Blue Ridge sectors, respectively.

Samples from four wells and one spring contained VOCs. Methyl tert-butyl ether (MTBE) occurred in samples from two wells, GWN-P1 and GWN-P15A, and one spring, GWN-P18. Chloroform was present in samples from spring GWN-P13A and well GWN-P7, both at levels considerably below the primary MCL (100 ppb total trihalomethanes). The chloroform occurrence in well GWN-P7 is one-time and probably resulted from the disinfection of the well following an overhaul. Trichloroethylene was present in the sample from well GWN-P1 and exceeded the primary MCL of 5 ppb. A follow-up sample confirmed the presence of excessive trichloroethylene. Well GWN-P16C yielded a sample containing vinyl chloride below the primary MCL. A follow-up sample contained no detectable vinyl chloride. The well had experienced an instance of vinyl chloride contamination in 1995. All of the stations producing samples with VOCs are located in or near built-up areas.

The fluoride content of the sample from spring GWN-P12A exceeded the secondary MCL of 2 ppm. The source of the fluoride in spring GWN-P12A is almost certainly natural.

### **3.10 VALLEY AND RIDGE UNCONFINED AQUIFERS**

Soil and residuum form low-yield unconfined aquifers across most of the Valley and Ridge Province of northwestern Georgia. Valley bottoms underlain by dolostones and limestones of the Cambro-Ordovician Knox Group are the locations of most higher-yielding wells and springs that are suitable for municipal supplies.

Five wells and four springs were used to monitor the water quality in the Valley and Ridge unconfined aquifers (Figure 3-18). Three of the wells and all four springs produced water from Knox Group carbonates. Of the remaining wells, well GWN-VR4 derives water from the Ordovician Chickamauga Group, while well GWN-VR6 taps the Cambrian Shady Dolomite.






Figure 3-15 Locations of Stations Monitoring the Piedmont/Blue Ridge Unconfined Aquifers



Nitrate/nitrite levels below the detection limit are assigned a value of 0.005 ppm. A missing bar indicates that samples were not collected for that year.

Figure 3-16 Nitrate/Nitrite Concentrations for Selected Wells in the Piedmont/Blue Ridge Unconfined Aquifer System: Piedmont Sector



Nitrate/nitrite levels below the detection limit are assigned a value of 0.007 ppm. A missing bar indicates that samples were not collected for that year.

Figure 3-17 Nitrate/Nitrite Concentrations for Selected Wells in the Piedmont/Blue Ridge Unconfined Aquifer System: Blue Ridge Sector







Figure 3-18 Locations of Stations Monitoring the Valley and Ridge Unconfined Aquifers

Sample pHs were mostly basic and ranged from 6.93 to 7.75. Conductivities ranged from 184 uS to 500 uS. All samples were tested for nitrate/nitrite and for VOCs.

Nitrate/nitrite ranged from undetected to 2.6 ppm as nitrogen. Figure 3-19 shows nitrite/nitrate levels for selected sampling stations in the Valley and Ridge aquifers. VOCs were present in samples from two stations. One of these, spring GWN-VR8, located near a commercial area, gave a sample containing a low level of MTBE. The spring has intermittently experienced contamination from motor fuel components in the past. The other, well GWN-VR6, gave a sample containing chloroform, 1,1-dichloroethylene, and tetrachloroethylene. The well is located in an industrial area and has, in the past, given samples contaminated with chlorinated aliphatic compounds. None of the volatile organic compounds exceeded the primary MCLs. Appendix Table A-11 presents the analytical summary for the wells and springs located in the Valley and Ridge unconfined aquifers.



A missing bar indicates that samples were not collected for that year.

Figure 3-19 Nitrate/Nitrite Concentrations for Selected Wells and Springs in the Valley and Ridge Unconfined Aquifers

## CHAPTER 4 SUMMARY AND CONCLUSIONS

EPD personnel collected 125 raw water samples from 114 wells and seven springs on the Ground-Water Monitoring Network during calendar year 2000 for volatile organic and limited inorganic analysis. These wells and springs monitor the water quality of nine aquifer systems in Georgia:

- Cretaceous aquifer system
- Providence aquifer system
- Clayton aquifer system
- Claiborne aquifer system
- Jacksonian aquifer system
- Floridan aquifer system
- Miocene aquifer system
- Piedmont/Blue Ridge unconfined aquifers
- Valley and Ridge unconfined aquifers

Comparisons of analyses of water samples collected during calendar year 2000 were made with analyses for the Ground-Water Monitoring Network dating back to 1984, permitting the recognition of temporal trends. Table 4-1 lists the contaminants and pollutants detected at the stations of the Ground-Water Monitoring Network during 2000. Although isolated water quality problems existed at specific localities, the quality of water from most of the Ground-Water Monitoring Network stations remains excellent.

Nitrate/nitrite is the most common substance(s) present in ground water in Georgia that can have adverse health effects. One well (GWN-MI15), a shallow domestic well tapping the Miocene aquifer system and located near a row crop field, yielded a water sample with a nitrite/nitrate concentration exceeding the primary MCL of 10 ppm as nitrogen (Table 4-1). (The owner of the well received notification about the excess nitrate/nitrite, and all well owners receive copies of the analytical results.) Samples from four other wells (GWN-J8, GWN-MI5, GWN-MI7, and GWN-MI9A) also had nitrate/nitrite levels that were elevated though not greater than the primary MCL. All four are shallow domestic-type wells. Well GWN-MI5 is located near a newly erected and occupied animal enclosure. The other three wells are located near row crop fields. A substantial increase in the nitrate/nitrite level for well GWN-MI9A occurred when the adjacent field, left fallow during 1999, was planted in corn during 2000. The increase may have resulted from fertilizer application. The nitrate/nitrite level declined between 1999 and 2000 for well GWN-CT7A when animals were removed from a nearby enclosure.

Spatial and temporal limitations of the Ground-Water Monitoring Network preclude the identification of the exact sources of the increasing levels of nitrogen compounds

Table 4-1. Pollution and Contamination Incidents, Calendar Year 2000.

Station	Contaminant/Pollutant	Primary MCL
GWN-K1	CHCl <sub>3</sub> =1.5ppb C <sub>2</sub> HCl <sub>3</sub> =0.64ppb	total trihalomethanes=100ppb C <sub>2</sub> HCl <sub>3</sub> =5ppb
GWN-K5	C <sub>2</sub> HCl <sub>3</sub> =0.88ppb/1.1ppb	C <sub>2</sub> HCl <sub>3</sub> =5ppb
GWN-K16	CHCl <sub>3</sub> =4.2ppb CHBrCl <sub>2</sub> =1.5ppb	total trihalomethanes=100ppb
GWN-J8	Be=4.3ppb	Be=4ppb
GWN-MI15	NO <sub>x</sub> =27ppm as N	NO <sub>x</sub> =10ppm as N
GWN-PA17	CHCl <sub>3</sub> =0.91ppb CHBrCl <sub>2</sub> =0.75ppb CHClBr <sub>2</sub> =0.61ppb	total trihalomethanes=100ppb
GWN-PA33A	CHCl <sub>3</sub> =2.1ppb CHBrCl <sub>2</sub> =0.79ppb	total trihalomethanes=100ppb
GWN-P1	C <sub>2</sub> HCl <sub>3</sub> =6.7ppb/6.0ppb/3.8ppb MTBE=8.0ppb/5.5ppb/5.5ppb	C <sub>2</sub> HCl <sub>3</sub> =5ppb (none)
GWN-P7	CHCl <sub>3</sub> =0.59ppb	total trihalomethanes=100ppb
GWN-P12A	F=3.8ppm Cl=12ppm SO <sub>4</sub> =65ppm	F=4ppm (none) (none)
GWN-P13A	CHCl <sub>3</sub> =0.76ppb	total trihalomethanes=100ppb
GWN-P15A	MTBE=0.78ppb	(none)
GWN-P16C	C <sub>2</sub> H <sub>3</sub> Cl=1.1ppb	C <sub>2</sub> H <sub>3</sub> Cl=2ppb
GWN-P18	MTBE=0.51ppb	(none)
GWN-VR6	CHCl <sub>3</sub> =0.56ppb 1,1- C <sub>2</sub> H <sub>2</sub> Cl <sub>2</sub> =1.7ppb C <sub>2</sub> Cl <sub>4</sub> =2.8ppb	total trihalomethanes=100ppb 1,1- C <sub>2</sub> H <sub>2</sub> Cl <sub>2</sub> =7ppb C <sub>2</sub> Cl <sub>4</sub> =5ppb
GWN-VR8	MTBE=1.2ppb	(none)

Notes:

NO<sub>x</sub> = Nitrate/Nitrite

MTBE = Methyl-tert-butyl Ether

A slash (/) separates values if multiple samples were taken from one station.

in some of Georgia's ground water. Nitrite/nitrate originates in ground water from direct sources and through oxidation of other forms of dissolved nitrogen, deriving from both natural and manmade sources. The most common sources of manmade dissolved nitrogen in Georgia usually consist of septic systems, agricultural wastes, and storage or application of fertilizers (Robertson, et. al., 1993). Dissolved nitrogen also is present in rainwater and can be derived from terrestrial vegetation and volatilization of fertilizers (Drever, 1988). The conversion of other nitrogen species to nitrate occurs in aerobic environments such as recharge areas. Anaerobic conditions in ground water, which commonly develop along the flow path of ground water, foster the denitrification process. However, the lack of denitrifying bacteria in ground water may inhibit this process (Freeze and Cherry, 1979).

Volatile organic compounds were detected in samples from thirteen stations. MTBE was detected in samples from two wells (GWN-P1 and GWN-P15A) and two springs (GWN-P18 and GWN-VR8). All four of these stations are located in or near built-up areas, and spring GWN-VR8 has experienced intermittent low-level pollution by motor fuel components in the past. Samples from six wells and one spring contained low levels of the trihalomethane, chloroform. For wells GWN-PA17 and GWN-PA33A, the chloroform probably originated from the reflux of chlorinated water down the well bores. The chlorine in the water then reacts with naturally occurring dissolved organic matter to form the chloroform. Overhauls of wells GWN-K16 and GWN-P7 were completed only a short time before sampling, and the chloroform probably arose during disinfection prior to placing them back in service. For spring GWN-P13A and wells GWN-K1 and GWN-VR6, the reason for the presence of chloroform is not clear. Samples from five wells were contaminated with chlorinated ethane and ethylene compounds, with the level of trichloroethylene for well GWN-P1 exceeding the primary MCL. Three of the wells (GWN-K1, GWN-K5, and GWN-VR6) are located in industrial settings. The remaining ones (GWN-P1 and GWN-P16C) are located in or near built-up areas.

Beryllium exceeded the primary MCL (4 ppb) in the sample from well GWN-J8, and fluoride exceeded the secondary MCL (2 ppm) in the sample from spring GWN-P12A. Well GWN-J8 is a domestic water well located in the Coastal Plain, which has intermittently yielded samples with excessive beryllium in the past. Spring GWN-P12A is located on the Piedmont and has previously given samples containing excessive fluoride. A sign placed near the spring advises against consuming the water. The sources of the beryllium and the fluoride are both almost certainly natural.

Field pH measurements taken during the year 2000 for various stations were compared with those taken during previous years for the same stations. The comparison found measurements taken from June into September of 2000 were consistently lower, by about a half unit to about a unit, than measurements taken at the same stations during previous years. The stations tested during this period cover several different aquifers. This variation may be related to the drought conditions occurring in Georgia since May 1998, or may be due to some type of systematic error. Possible sources of error

include a subtle instrument malfunction, off-quality calibration standards, or off-quality distilled water for dissolving the standards.

A manufacturer's technical representative re-emphasized that almost any type of malfunction involving the model of instrument used for the pH measurements would cause the display of some sort of error message. Instrument malfunction would then be unlikely as an error source. Calibration standards are prepared from packages of powders each sufficient for making up 50 milliliters of standard solution (the amount usually used for a single calibration). The (inadvertent) use of some packages of off-quality standard powders would likely have caused problems before June and after September. If the low pH measurements were due to systematic error, off-quality distilled water would be the most likely source. The above-mentioned technical representative suggested a quick method for testing the quality of distilled water.

## CHAPTER 5 LIST OF REFERENCES

- Brooks, R., Clarke, J.S., and Faye, R.E., 1985, Hydrology of the Gordon Aquifer System of East-Central Georgia: Georgia Geologic Survey Information Circular 75, 41 p., 2 pl.
- Clarke, J.S., Faye, R.E., and Brooks, R., 1983, Hydrogeology of the Providence Aquifer of Southwest Georgia: Georgia Geologic Survey Hydrologic Atlas 11, 5 pl.
- Clarke, J.S., Faye, R.E., and Brooks, R., 1984, Hydrogeology of the Clayton Aquifer of Southwest Georgia: Georgia Geologic Survey Hydrologic Atlas 13, 6 pl.
- Clarke, J.S., Brooks, R., and Faye, R.E., 1985, Hydrogeology of the Dublin and Midville Aquifer Systems of East Central Georgia: Georgia Geologic Survey Information Circular 74, 62 p., 2 pl.
- Clarke, J.S., Hacke, C.M., and Peck, M.F., 1990, Geology and Ground-Water Resources of the Coastal Area of Georgia: Georgia Geologic Survey Bulletin 113, 116 p., 12 pl.
- Clarke, J.S., Falls, W.F., Edwards, L.E., Frederiksen, N.O., Bybell, L.M., Gibson, T.G., Gohn, G.S., and Fleming, F., 1996, Hydrogeologic Data and Aquifer Interconnection in a Multi-Aquifer System in Coastal Plain Sediments Near Millhaven, Screven County, Georgia: Georgia Geologic Survey Information Circular 99, 49p. 1pl.
- Crews, P.A., and Huddleston, P.F., 1984, Geologic Sections of the Principal Artesian Aquifer System, in Hydrogeologic Evaluation for Underground Injection Control in the Coastal Plain of Georgia, R. Arora, ed.: Georgia Geologic Survey Hydrologic Atlas 10, 41pl.
- Davis, K.R., Donahue, J.C., Hutcheson, R.H., and Waldrop, D.L., 1988, Most Significant Ground-Water Recharge Areas of Georgia: Georgia Geologic Survey Hydrologic Atlas 18, 1 pl.
- Drever, J. I., 1988, The Geochemistry of Natural Waters: Prentice-Hall, Englewood Cliffs, New Jersey, 437 p.
- EPD, 1991, A Ground-Water Management Plan for Georgia: Georgia Geologic Survey Circular 11 (1991 edition).
- EPD, 1997, Rules for Safe Drinking Water, Section 391-3-5, Rules of the Georgia Department of Natural Resources Environmental Protection Division, p. 601-691.



- EPD, 1998, A Ground-Water Management Plan for Georgia: Georgia Geologic Survey Circular 11 (1998 edition).
- Freeze, R.A., and Cherry, J.A., 1979, Groundwater: Prentice-Hall, Englewood Cliffs, New Jersey, 604 p.
- Gorday, L.L., Lineback, J.A., Long, A.F., McLemore, W.H., 1997, A Digital Model Approach to Water-Supply Management of the Claiborne, Clayton, and Providence Aquifers of Southwestern Georgia: Georgia Geologic Survey Bulletin 118, 31 p., Appendix, Supplements I and II.
- Hayes, L.R., Maslia, M.L., and Meeks, W.C., 1983, Hydrology and Model Evaluation of the Principal Artesian Aquifer, Dougherty Plain, Southwest Georgia: Georgia Geologic Survey Bulletin 97, 93p.
- Hicks, D.W., Krause, R.E., and Clarke, J.S., 1981, Geohydrology of the Albany Area, Georgia: Georgia Geologic Survey Information Circular 57, 31 p.
- Hetrick, J.H., 1990, Geologic Atlas of the Fort Valley Area: Georgia Geologic Survey Geologic Atlas 7, 2 pl.
- Hetrick, J.H., 1992, A Geologic Atlas of the Wrens-Augusta Area: Georgia Geologic Survey Geologic Atlas 8, 3 pl.
- Huddleston, P.F. and Summerour, J.H., 1996, The Lithostratigraphic Framework of the Uppermost Cretaceous and Lower Tertiary of Eastern Burke County, Georgia: Georgia Geologic Survey Bulletin 127, 94 p., 1 pl.
- Joiner, C.N., Reynolds, M.S., Stayton, W.L., and Boucher, F.G., 1988, Ground-Water Data for Georgia, 1987: United States Geological Survey Open-File Report 88-323, 172 p.
- Kellam, M.F., and Gorday, L.L., 1990, Hydrogeology of the Gulf Trough-Apalachicola Embayment Area, Georgia: Georgia Geologic Survey Bulletin 94, 74 p., 15 pl.
- Krause, R.E., 1979, Geohydrology of Brooks, Lowndes, and Western Echols Counties, Georgia: United States Geological Survey Water-Resources Investigations 78-117, 48 p., 8 pl.
- Long, A.F., 1989, Hydrogeology of the Clayton and Claiborne Aquifer Systems: Georgia Geologic Survey Hydrologic Atlas 19, 6 pl.
- McFadden, S.S., and Perriello, P.D., 1983, Hydrogeology of the Clayton and Claiborne Aquifers in Southwestern Georgia: Georgia Geologic Survey Information Circular 55, 59 p., 2 pl.

- Pollard, L.D., and Vorhis, R.C., 1980, The Geohydrology of the Cretaceous Aquifer System in Georgia: Georgia Geologic Survey Hydrologic Atlas 3, 5 pl.
- Robertson, S.J., Shellenberger, D.L., York, G.M., Clark, M.G., Eppihimer, R.M., Lineback, J.A., 1993, Sampling for Nitrate Concentrations in North Georgia's Ground Water: 1993 Georgia Water Resources Conference 364-365, 1 p.
- Shellenberger, D.L., Barget, R.G., Lineback, J.A., and Shapiro, E.A., 1996, Nitrate in Georgia's Ground Water: Georgia Geologic Survey Project Report 25, 12 p., 1 pl.
- Stuart, M.A., Rich, F.J., and Bishop, G.A., 1995, Survey of Nitrate Contamination in Shallow Domestic Drinking Water Wells in the Inner Coastal Plain of Georgia: Ground Water, Vol. 33, No. 2, p. 284-290.
- Summerour, J.H., Shapiro, E.A., Lineback, J.A., Huddleston, P.F., and Hughes, A.C., 1994, An Investigation of Tritium in the Gordon and Other Aquifers in Burke County, Georgia: Georgia Geologic Survey Information Circular 95, 93 p.
- Tolford, B., 1999, Pesticide Monitoring Network 1998-1999: Georgia Geologic Survey Project Report 40, 60 p.
- Tuohy, M.A., 1984, Isopach Map of the Claiborne Aquifer, in Hydrogeologic Evaluation for Underground Injection Control in the Coastal Plain of Georgia, R. Arora, ed.: Georgia Geologic Survey Hydrologic Atlas 10.
- Vincent, R.H., 1982, Geohydrology of the Jacksonian Aquifer in Central and East Central Georgia: Georgia: Geologic Survey Hydrologic Atlas 8, 3 pl.
- Watson, T., 1982, Aquifer Potential of the Shallow Sediments of the Coastal Area of Georgia: Proceedings, Second Symposium on the Geology of the Southeastern Coastal Plain, Arden, D.D., Beck, B.F., Morrow, E., eds., Georgia Geologic Survey Information Circular 53, p. 183-194.

## **APPENDIX**

Laboratory Data

## LABORATORY DATA

The standard testing regimen for all samples collected for the Ground-Water Monitoring Network consisted of laboratory analyses for volatile organic compounds and nitrate/nitrite and of field measurements of pH and electrical conductivity. Tests were carried out at two stations for additional substances, which are subject to primary MCL's and had histories of occurrence at these stations.

EPA has set forth a series of (serially numbered) analytical methods officially recognized as suitable for environmental purposes. The EPD laboratory cites EPA method numbers along with analysis results, and Tables A-1 and A-2 list the method numbers appropriate to the various analytes.

Tables A-3 through A-11 regularly list results for the following parameters: pH, conductivity, nitrate/nitrite, trihalomethanes, MTBE, benzene, toluene, ethyl benzene, and total xylenes. Other VOC's are listed if detected. Results for one station also list beryllium; and, results for another station substances amenable to EPA method 300.0 -- fluoride, chloride, and sulfate. The abbreviation "ppm", where used in a nitrate/nitrite entry in these tables, is understood to mean milligrams per liter as nitrogen.

For this appendix, the following abbreviations are used:

su	= standard units
mg/L	= milligrams per liter (parts per million)
ppm	= parts per million
mg/L as N	= milligrams per liter (parts per million), as nitrogen
ug/L	= micrograms per liter (parts per billion)
ppb	= parts per billion
uS/cm	= microsiemens/centimeter
nd	= not detected
--	= not analyzed

Note:

The detection limit for the same substance can vary among different laboratories and can vary for a single laboratory if a sample is diluted to lower the concentration of interfering substances, or if the array of standards used to develop the detection limit is revised.

Table A-1. Standard Water Quality Analyses: Anions, Volatile Organic Compounds, and Other Parameters.

<b>ANIONS</b>			
<b>Parameter</b>	<b>Test Method</b>	<b>Method Detection Limit</b>	<b>Primary Maximum Contaminant Level</b>
Nitrate/Nitrite (NO <sub>x</sub> )	EPA 353.2	0.02 mg/L as N	10 mg/L as N

<b>VOLATILE ORGANIC COMPOUNDS</b>			
<b>Parameter</b>	<b>Type of Test</b>	<b>Practical Quantitation Limit</b>	<b>Primary Maximum Contaminant Level</b>
Vinyl Chloride	EPA 524.2	0.5 ug/L	2.0 ug/L
1,1-Dichloroethylene	EPA 524.2	0.5 ug/L	7.0 ug/L
Dichloromethane	EPA 524.2	0.5 ug/L	5.0 ug/L
Trans-1,2-Dichloroethylene	EPA 524.2	0.5 ug/L	100 ug/L
Cis-1,2-Dichloroethylene	EPA 524.2	0.5 ug/L	70.0 ug/L
1,1,1-Trichloroethane	EPA 524.2	0.5 ug/L	200 ug/L
Carbon Tetrachloride	EPA 524.2	0.5 ug/L	5.0 ug/L
Benzene	EPA 524.2	0.5 ug/L	5.0 ug/L
1,2-Dichloroethane	EPA 524.2	0.5 ug/L	5.0 ug/L
Trichloroethylene	EPA 524.2	0.5 ug/L	5.0 ug/L
1,2-Dichloropropane	EPA 524.2	0.5 ug/L	5.0 ug/L
Toluene	EPA 524.2	0.5 ug/L	1000 ug/L
1,1,2-Trichloroethane	EPA 524.2	0.5 ug/L	5.0 ug/L
Tetrachloroethylene	EPA 524.2	0.5 ug/L	5.0 ug/L
Chlorobenzene	EPA 524.2	0.5 ug/L	100 ug/L
Ethylbenzene	EPA 524.2	0.5 ug/L	700 ug/L
Total Xylenes	EPA 524.2	0.5 ug/L	10,000 ug/L

Table A-1 (Continued). Standard Water Quality Analyses: Anions, Volatile Organic Compounds, and Other Parameters.

<b>VOLATILE ORGANIC COMPOUNDS</b>			
<b>Parameter</b>	<b>Type of Test</b>	<b>Practical Quantitation Limit</b>	<b>Primary Maximum Contaminant Level</b>
Styrene	EPA 524.2	0.5 ug/L	100 ug/L
1,4-Dichlorobenzene (P)	EPA 524.2	0.5 ug/L	75.0 ug/L
1,2-Dichlorobenzene (O)	EPA 524.2	0.5 ug/L	600 ug/L
1,2,4-Trichlorobenzene	EPA 524.2	0.5 ug/L	70.0 ug/L
Dichlorodifluoromethane	EPA 524.2	0.5 ug/L	None
Chloromethane	EPA 524.2	0.5 ug/L	None
Bromomethane	EPA 524.2	0.5 ug/L	None
Chloroethane	EPA 524.2	0.5 ug/L	None
Trichlorofluoromethane	EPA 524.2	0.5 ug/L	None
1,1-Dichloroethane	EPA 524.2	0.5 ug/L	None
2,2-Dichloropropane	EPA 524.2	0.5 ug/L	None
Bromochloromethane	EPA 524.2	0.5 ug/L	None
Chloroform	EPA 524.2	0.5 ug/L	100 ug/L*
1,1-Dichloropropylene	EPA 524.2	0.5 ug/L	None
Dibromomethane	EPA 524.2	0.5 ug/L	None
Bromodichloromethane	EPA 524.2	0.5 ug/L	100 ug/L*
Cis-1,3-Dichloropropylene	EPA 524.2	0.5 ug/L	None
Trans-1,3-Dichloropropylene	EPA 524.2	0.5 ug/L	None
1,3-Dichloropropane	EPA 524.2	0.5 ug/L	None
Dibromochloromethane	EPA 524.2	0.5 ug/L	100 ug/L*

Table A-1 (Continued). Standard Water Quality Analyses: Anions, Volatile Organic Compounds, and Other Parameters.

<b>VOLATILE ORGANIC COMPOUNDS</b>			
<b>Parameter</b>	<b>Type of Test</b>	<b>Practical Quantitation Limit</b>	<b>Primary Maximum Contaminant Level</b>
1,2-Dibromoethane	EPA 524.2	0.5 ug/L	None
1,1,1,2-Tetrachloroethane	EPA 524.2	0.5 ug/L	None
Bromoform	EPA 524.2	0.5 ug/L	100 ug/L*
Isopropylbenzene	EPA 524.2	0.5 ug/L	None
1,1,2,2-Tetrachloroethane	EPA 524.2	0.5 ug/L	None
Bromobenzene	EPA 524.2	0.5 ug/L	None
1,2,3-Trichloropropane	EPA 524.2	0.5 ug/L	None
N-Propylbenzene	EPA 524.2	0.5 ug/L	None
2-Chlorotoluene	EPA 524.2	0.5 ug/L	None
1,3,5-Trimethylbenzene	EPA 524.2	0.5 ug/L	None
4-Chlorotoluene	EPA 524.2	0.5 ug/L	None
Tert-Butylbenzene	EPA 524.2	0.5 ug/L	None
1,2,4-Trimethylbenzene	EPA 524.2	0.5 ug/L	None
Sec-Butylbenzene	EPA 524.2	0.5 ug/L	None
P-Isopropyltoluene	EPA 524.2	0.5 ug/L	None
1,3-Dichlorobenzene (M)	EPA 524.2	0.5 ug/L	None
N-Butylbenzene	EPA 524.2	0.5 ug/L	None
1,2-Dibromo-3-Chloropropane	EPA 524.2	0.5 ug/L	0.2 ug/L
Hexachlorobutadiene	EPA 524.2	0.5 ug/L	None
Naphthalene	EPA 524.2	0.5 ug/L	None

Table A-1 (Continued). Standard Water Quality Analyses: Anions, Volatile Organic Compounds, and Other Parameters.

<b>VOLATILE ORGANIC COMPOUNDS</b>			
<b>Parameter</b>	<b>Type of Test</b>	<b>Practical Quantitation Limit</b>	<b>Primary Maximum Contaminant Level</b>
1,2,3-Trichlorobenzene	EPA 524.2	0.5 ug/L	None
Methyl Tert-butyl Ether	EPA 524.2	0.5 ug/L	None

<b>OTHER PARAMETERS**</b>		
<b>Parameter</b>	<b>Units</b>	<b>Maximum Contaminant Level</b>
pH	0.01 SU	None
Conductivity	1.0 uS	None

Notes:

Detection limits for analyses are Practical Quantitation Limits (PQLs) except for the NO<sub>x</sub> test, for which a Method Detection Limit (MDL) is used.

Primary MCL's from Georgia Rules for Safe Drinking Water, October 1997 edition (EPD, 1997).

\* Indicates a trihalomethane compound. The primary MCL for total trihalo- methanes is 100 ug/L.

\*\*pH and conductivity are measured in the field (see Chapter 2).



Table A-2. Optional Water Quality Analyses: Metals and Anions.

<b>METALS</b>			
<b>Parameter</b>	<b>Test Method</b>	<b>Practical Quantitation Limit</b>	<b>Max. Contaminant Level</b>
Beryllium (Be)	EPA 210.2	2 ug/L	4 ug/L <sub>1</sub>

<b>ANIONS</b>			
<b>Parameter</b>	<b>Test Method</b>	<b>Method Detection Limit</b>	<b>Max. Contaminant Level</b>
Bromide (Br)	EPA 300.0	0.1 mg/L	None
Chloride (Cl)	EPA 300.0	0.1 mg/L	250 mg/L <sub>2</sub>
Sulfate (SO <sub>4</sub> )	EPA 300.0	2.0 mg/L	250 mg/L <sub>2</sub>
Fluoride (F)	EPA 300.0	0.1 mg/L	4.0 mg/L <sub>1</sub> , 2.0 mg/L <sub>2</sub>

Notes:

Detection limits for anions analyses are Method Detection Limits (MDLs). The detection limit for beryllium is a Practical Quantitation Limit (PQL).

MCL's from Georgia Rules for Safe Drinking Water, October 1997 edition (EPD, 1997):

<sub>1</sub>=Primary Maximum Contaminant Level (MCL).

<sub>2</sub>=Secondary MCL.

Table A-3. 2000 Ground-Water Quality Analyses of the Cretaceous Aquifer System.

GWN-K1			
Well Name:	Englehard Kaolin Company #2		
County:	Wilkinson		
Date Sampled:	03/09/2000		
Nitrate/Nitrite	0.68	ppm	
pH	4.21	su	
conductivity	231	uS/cm	
Trihalomethanes: chloroform	1.5	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	0.64	ppb	
Other			

GWN-K2A			
Well Name:	Irwinton #303		
County:	Wilkinson		
Date Sampled:	03/09/2000		
Nitrate/Nitrite	0.07	ppm	
pH	5.80	su	
conductivity	96	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K3			
Well Name:	Sandersville #7B		
County:	Washington		
Date Sampled:	03/09/2000		
Nitrate/Nitrite	0.04	ppm	
pH	5.93	su	
conductivity	110	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K5			
Well Name:	Richmond County #101		
County:	Richmond		
Date Sampled:	05/03/2000		
Nitrate/Nitrite	0.95	ppm	
pH	4.27	su	
conductivity	21	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	0.88	ppb	
Other			

GWN-K5			
Well Name:	Richmond County #101		
County:	Richmond		
Date Sampled:	09/19/2000		
Nitrate/Nitrite	0.95	ppm	
pH	--	su	
conductivity	20	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	1.1	ppb	
Other			

GWN-K6			
Well Name:	Huber #6		
County:	Twiggs		
Date Sampled:	06/22/2000		
Nitrate/Nitrite	0.031	ppm	
pH	4.88	su	
conductivity	31	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-3 (Continued). 2000 Ground-Water Quality Analyses of the Cretaceous Aquifer System.

GWN-K7			
Well Name:	Jones County #4		
County:	Jones		
Date Sampled:	06/22/2000		
Nitrate/Nitrite	0.13	ppm	
pH	4.26	su	
conductivity	14	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K8			
Well Name:	Mohawk Industries #3		
County:	Laurens		
Date Sampled:	12/12/2000		
Nitrate/Nitrite	0.043	ppm	
pH	6.76	su	
conductivity	180	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K9			
Well Name:	Marshallville #1		
County:	Macon		
Date Sampled:	02/24/2000		
Nitrate/Nitrite	nd	ppm	
pH	4.32	su	
conductivity	50	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K10B			
Well Name:	Fort Valley #6		
County:	Peach		
Date Sampled:	12/12/2000		
Nitrate/Nitrite	0.62	ppm	
pH	4.92	su	
conductivity	13	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K11A			
Well Name:	Warner Robins #2		
County:	Houston		
Date Sampled:	09/21/2000		
Nitrate/Nitrite	0.82	ppm	
pH	4.79	su	
conductivity	20	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K12			
Well Name:	Perry/Holiday Inn Well		
County:	Houston		
Date Sampled:	09/21/2000		
Nitrate/Nitrite	nd	ppm	
pH	4.21	su	
conductivity	39	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-3 (Continued). 2000 Ground-Water Quality Analyses of the Cretaceous Aquifer System.

GWN-K13			
Well Name:	Omaha #1		
County:	Stewart		
Date Sampled:	04/05/2000		
Nitrate/Nitrite	nd	ppm	
pH	8.92	su	
conductivity	244	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K15A			
Well Name:	Georgetown #3		
County:	Quitman		
Date Sampled:	04/05/2000		
Nitrate/Nitrite	nd	ppm	
pH	9.22	su	
conductivity	372	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K16			
Well Name:	Pactiv, Inc. North Well		
County:	Bibb		
Date Sampled:	12/14/2000		
Nitrate/Nitrite	0.52	ppm	
pH	5.98	su	
conductivity	39	uS/cm	
Trihalomethanes: chloroform	4.2	ppb	
bromodichloromethane	1.5		
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K18A			
Well Name:	Buena Vista #6		
County:	Marion		
Date Sampled:	04/04/2000		
Nitrate/Nitrite	0.15	ppm	
pH	4.49	su	
conductivity	23	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K19			
Well Name:	Hephzibah/Murphy St. Well		
County:	Richmond		
Date Sampled:	05/03/2000		
Nitrate/Nitrite	0.10	ppm	
pH	4.59	su	
conductivity	16	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-K20			
Well Name:	Plains #7		
County:	Sumter		
Date Sampled:	04/04/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.39	su	
conductivity	136	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-4. 2000 Ground-Water Quality Analyses of the Providence Aquifer System.

GWN-PD2B			
Well Name:	Preston #4		
County:	Webster		
Date Sampled:	04/04/2000		
Nitrate/Nitrite	1.1	ppm	
pH	5.45	su	
conductivity	48	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PD3			
Well Name:	Fort Gaines #2		
County:	Clay		
Date Sampled:	04/05/2000		
Nitrate/Nitrite	nd	ppm	
pH	8.38	su	
conductivity	389	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-5. 2000 Ground-Water Quality Analyses of the Clayton Aquifer System.

GWN-CT2A			
Well Name:	Burton Thomas house well		
County:	Sumter		
Date Sampled:	08/10/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.33	su	
conductivity	239	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CT3			
Well Name:	Dawson/ Crawford St. Well		
County:	Terrell		
Date Sampled:	08/10/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.33	su	
conductivity	189	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CT5A			
Well Name:	Cuthbert #3		
County:	Randolph		
Date Sampled:	08/10/2000		
Nitrate/Nitrite	0.026	ppm	
pH	7.29	su	
conductivity	183	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CT7A			
Well Name:	St. John farm well		
County:	Sumter		
Date Sampled:	10/12/2000		
Nitrate/Nitrite	3.6	ppm	
pH	4.64	su	
conductivity	48	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CT8			
Well Name:	Weathersby house well		
County:	Schley		
Date Sampled:	10/12/2000		
Nitrate/Nitrite	0.71	ppm	
pH	4.94	su	
conductivity	22	uS	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-6. 2000 Ground-Water Quality Analyses of the Claiborne Aquifer System.

GWN-CL2			
Well Name:	Unadilla #3		
County:	Dooly		
Date Sampled:	02/24/2000		
Nitrate/Nitrite	0.25	ppm	
pH	7.42	su	
conductivity	209	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CL4A			
Well Name:	Plains #5		
County:	Sumter		
Date Sampled:	04/04/2000		
Nitrate/Nitrite	3.1	ppm	
pH	4.28	su	
conductivity	60	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CL6			
Well Name:	LTV/ Copperweld Central Supply Well		
County:	Early		
Date Sampled:	04/05/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.64	su	
conductivity	315	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CL8			
Well Name:	Flint River Nursery Office Well		
County:	Dooly		
Date Sampled:	02/24/2000		
Nitrate/Nitrite	nd	ppm	
pH	6.18	su	
conductivity	93	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-CL9			
Well Name:	Newton #3		
County:	Baker		
Date Sampled:	10/11/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.90	su	
conductivity	204	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-7. 2000 Ground-Water Quality Analyses of the Jacksonian Aquifer System.

GWN-J1B			
Well Name:	Quick house well		
County:	Burke		
Date Sampled:	11/16/2000		
Nitrate/Nitrite	2.6	ppm	
pH	7.25	su	
conductivity	219	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J2B			
Well Name:	Oakwood Village Mob. Home Park #3		
County:	Burke		
Date Sampled:	11/16/2000		
Nitrate/Nitrite	0.75	ppm	
pH	7.56	su	
conductivity	185	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J3			
Well Name:	J.W. Black house well		
County:	Emanuel		
Date Sampled:	08/24/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.41	su	
conductivity	141	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J4			
Well Name:	Wrightsville #4		
County:	Johnson		
Date Sampled:	03/08/2000		
Nitrate/Nitrite	0.18	ppm	
Ph	7.55	su	
conductivity	239	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J5			
Well Name:	Cochran #3		
County:	Bleckley		
Date Sampled:	02/24/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.37	su	
conductivity	356	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J6			
Well Name:	Wrens #3		
County:	Jefferson		
Date Sampled:	05/03/2000		
Nitrate/Nitrite	nd	ppm	
pH	6.46	su	
conductivity	150	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			



Table A-7 (Continued). 2000 Ground-Water Quality Analyses of the Jacksonian Aquifer System.

GWN-J7			
Well Name:	Templeton livestock well		
County:	Burke		
Date Sampled:	11/16/2000		
Nitrate/Nitrite	3.0	ppm	
pH	4.93	su	
conductivity	45	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-J8			
Well Name:	Kahn house well		
County:	Jefferson		
Date Sampled:	05/03/2000		
Nitrate/Nitrite	8.5	ppm	
pH	4.61	su	
conductivity	88	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other: Be	4.3	ppb	

Table A-8. 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA1			
Well Name:	Thunderbolt #1		
County:	Chatham		
Date Sampled:	09/20/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.25	su	
conductivity	236	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA2			
Well Name:	Savannah #13		
County:	Chatham		
Date Sampled:	09/20/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.31	su	
conductivity	186	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA3			
Well Name:	Grist Equipment Co. shop well		
County:	Chatham		
Date Sampled:	09/20/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.22	su	
conductivity	176	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA4			
Well Name:	Tybee #1		
County:	Chatham		
Date Sampled:	09/20/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.33	su	
conductivity	677	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA5A			
Well Name:	Interstate Paper #2		
County:	Liberty		
Date Sampled:	04/19/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.72	su	
conductivity	362	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA6			
Well Name:	Hinesville #5		
County:	Liberty		
Date Sampled:	04/19/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.74	su	
conductivity	324	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA7			
Well Name:	Darien New South Well		
County:	McIntosh		
Date Sampled:	04/19/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.52	su	
conductivity	680	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA8			
Well Name:	ITT Rayonnier #4D		
County:	Wayne		
Date Sampled:	04/19/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.63	su	
conductivity	419	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA9C			
Well Name:	Miller Ball Park TW 25		
County:	Glynn		
Date Sampled:	04/19/2000		
Nitrate/Nitrite	0.02	ppm	
pH	7.99	su	
conductivity	543	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA10B			
Well Name:	Gilman Paper #11		
County:	Camden		
Date Sampled:	04/18/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.27	su	
conductivity	1279	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other:		ppb	

GWN-PA11			
Well Name:	St. Marys #2		
County:	Camden		
Date Sampled:	04/18/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.31	su	
conductivity	795	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA12			
Well Name:	Folkston #3		
County:	Charlton		
Date Sampled:	04/18/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.30	su	
conductivity	749	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA13			
Well Name:	Waycross #3		
County:	Ware		
Date Sampled:	04/18/2000		
Nitrate/Nitrite	nd		ppm
pH	7.50		su
conductivity	444		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-PA14			
Well Name:	Statesboro #7		
County:	Bulloch		
Date Sampled:	05/04/2000		
Nitrate/Nitrite	0.02		ppm
pH	7.53		su
conductivity	234		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-PA15			
Well Name:	King Finishing Division Fire Well		
County:	Screven		
Date Sampled:	05/04/2000		
Nitrate/Nitrite	nd		ppm
pH	7.80		su
conductivity	251		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-PA16			
Well Name:	Millen #1		
County:	Jenkins		
Date Sampled:	03/08/2000		
Nitrate/Nitrite	nd		ppm
pH	7.58		su
conductivity	278		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-PA17			
Well Name:	Swainsboro #7		
County:	Emanuel		
Date Sampled:	08/24/2000		
Nitrate/Nitrite	0.035		ppm
pH	7.16		su
conductivity	297		uS/cm
Trihalomethanes: chloroform	0.91		ppb
bromodichloromethane	0.75		
chlorodibromomethane	0.61		
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-PA18			
Well Name:	Metter #2		
County:	Candler		
Date Sampled:	03/08/2000		
Nitrate/Nitrite	nd		ppm
pH	7.87		su
conductivity	207		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA19			
Well Name:	Douglas #4		
County:	Coffee		
Date Sampled:	04/20/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.50	su	
conductivity	415	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA20			
Well Name:	Lakeland #2		
County:	Lanier		
Date Sampled:	01/26/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.73	su	
conductivity	374	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA21A			
Well Name:	Valdosta New #4		
County:	Lowndes		
Date Sampled:	01/26/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.95	su	
conductivity	239	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA22			
Well Name:	Thomasville #6		
County:	Thomas		
Date Sampled:	06/08/2000		
Nitrate/Nitrite	0.14	ppm	
pH	7.30	su	
conductivity	404	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA23			
Well Name:	Cairo #8		
County:	Grady		
Date Sampled:	02/23/2000		
Nitrate/Nitrite	0.05	ppm	
pH	7.75	su	
conductivity	374	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA24			
Well Name:	Bainbridge #1		
County:	Decatur		
Date Sampled:	02/16/2000		
Nitrate/Nitrite	2.0	ppm	
pH	7.72	su	
conductivity	218	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA25			
Well Name:	Donalsonville /7th St. Well		
County:	Seminole		
Date Sampled:	02/16/2000		
Nitrate/Nitrite	1.6	ppm	
pH	7.40	su	
conductivity	280	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA26			
Well Name:	Colquitt #3		
County:	Miller		
Date Sampled:	02/16/2000		
Nitrate/Nitrite	2.1	ppm	
pH	7.58	su	
conductivity	224	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA27			
Well Name:	Camilla/Industrial Park Well		
County:	Mitchell		
Date Sampled:	06/07/2000		
Nitrate/Nitrite	0.53	ppm	
pH	7.23	su	
conductivity	228	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA28			
Well Name:	Moultrie #1		
County:	Colquitt		
Date Sampled:	02/17/2000		
Nitrate/Nitrite	0.02	ppm	
pH	7.85	su	
conductivity	484	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA29			
Well Name:	Adel #6		
County:	Cook		
Date Sampled:	01/26/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.72	su	
conductivity	360	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA30A			
Well Name:	Amoco/Nashville Mills #1		
County:	Berrien		
Date Sampled:	02/17/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.75	su	
conductivity	357	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA31			
Well Name:	Tifton #6		
County:	Tift		
Date Sampled:	02/17/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.56	su	
conductivity	273	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA32			
Well Name:	Ocilla #3		
County:	Irwin		
Date Sampled:	01/27/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.78	su	
conductivity	180	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA33A			
Well Name:	Fitzgerald #G		
County:	Ben Hill		
Date Sampled:	01/27/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.78	su	
conductivity	144	uS/cm	
Trihalomethanes: chloroform	2.1	ppb	
bromodichloromethane	0.79		
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA34			
Well Name:	McRae #2 (Telfair Ave.)		
County:	Telfair		
Date Sampled:	12/06/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.73	su	
conductivity	339	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA36			
Well Name:	Vidalia #1		
County:	Toombs		
Date Sampled:	08/24/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.57	su	
conductivity	178	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA38			
Well Name:	Eastman #4		
County:	Dodge		
Date Sampled:	12/06/2000		
Nitrate/Nitrite	0.24	ppm	
pH	7.83	su	
conductivity	232	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA39			
Well Name:	Sylvester #1		
County:	Worth		
Date Sampled:	01/27/2000		
Nitrate/Nitrite	0.04	ppm	
pH	7.42	su	
conductivity	305	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA40			
Well Name:	Merck #8		
County:	Dougherty		
Date Sampled:	06/07/2000		
Nitrate/Nitrite	1.6	ppm	
pH	7.11	su	
conductivity	287	uS/cm	
Trihalomethanes:	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA43A			
Well Name:	Pineland Fish Farm office well		
County:	Baker		
Date Sampled:	06/07/2000		
Nitrate/Nitrite	3.1	ppm	
pH	7.13	su	
conductivity	290	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA44			
Well Name:	Sycamore #2		
County:	Turner		
Date Sampled:	07/27/2000		
Nitrate/Nitrite	0.18	ppm	
pH	7.31	su	
conductivity	143	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA45A			
Well Name:	Abbeville #1		
County:	Wilcox		
Date Sampled:	07/27/2000		
Nitrate/Nitrite	0.45	ppm	
pH	7.08	su	
conductivity	193	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA46B			
Well Name:	Wenona Mobile Home Park Well		
County:	Crisp		
Date Sampled:	11/02/2000		
Nitrate/Nitrite	3.2	ppm	
pH	7.71	su	
conductivity	188	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			



Table A-8 (Continued). 2000 Ground-Water Quality Analyses of the Floridan Aquifer System.

GWN-PA49			
Well Name:	Harmony Church Well		
County:	Dooly		
Date Sampled:	06/22/2000		
Nitrate/Nitrite	1.6	ppm	
pH	7.32	su	
conductivity	201	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA50			
Well Name:	Reynolds house well		
County:	Laurens		
Date Sampled:	11/15/2000		
Nitrate/Nitrite	1.2	ppm	
pH	7.52	su	
conductivity	294	uS/cm	
Trihalomethanes:	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA51			
Well Name:	Adams house well		
County:	Mitchell		
Date Sampled:	02/16/2000		
Nitrate/Nitrite	1.0	ppm	
pH	7.55	su	
conductivity	220	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA52			
Well Name:	Simmons house well		
County:	Mitchell		
Date Sampled:	02/23/2000		
Nitrate/Nitrite	3.3	ppm	
pH	7.78	su	
conductivity	224	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA53			
Well Name:	Cato house well		
County:	Decatur		
Date Sampled:	02/23/2000		
Nitrate/Nitrite	4.4	ppm	
pH	7.78	su	
conductivity	228	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-PA55			
Well Name:	Holland house well		
County:	Burke		
Date Sampled:	11/15/2000		
Nitrate/Nitrite	0.052	ppm	
pH	7.64	su	
conductivity	194	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-9. 2000 Ground-Water Quality Analyses of the Miocene Aquifer System.

GWN-MI1			
Well Name:	McMillan house well		
County:	Cook		
Date Sampled:	06/08/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.28	su	
conductivity	239	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-MI2A			
Well Name:	S. Boutwell house well		
County:	Lowndes		
Date Sampled:	06/08/2000		
Nitrate/Nitrite	2.1	ppm	
pH	3.80	su	
conductivity	57	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-MI5			
Well Name:	Carter house well		
County:	Appling		
Date Sampled:	11/14/2000		
Nitrate/Nitrite	6.1	ppm	
pH	5.15	su	
conductivity	60	uS	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-MI7			
Well Name:	Chaudoin house well		
County:	Irwin		
Date Sampled:	7/27/2000		
Nitrate/Nitrite	7.0	ppm	
pH	3.81	su	
conductivity	77	uS	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-MI9A			
Well Name:	Murphy garden well		
County:	Thomas		
Date Sampled:	10/11/2000		
Nitrate/Nitrite	9.6	ppm	
pH	6.27	su	
conductivity	114	uS	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-MI10B			
Well Name:	Calhoun house well		
County:	Colquitt		
Date Sampled:	12/13/2000		
Nitrate/Nitrite	nd	ppm	
pH	6.62	su	
conductivity	98	uS	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

Table A-9 (Continued). 2000 Ground-Water Quality Analyses of the Miocene Aquifer System.

GWN-MI15			
Well Name:	Aldrich house well		
County:	Bulloch		
Date Sampled:	05/04/2000		
Nitrate/Nitrite	27		ppm
pH	4.37		su
conductivity	157		uS
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

Table A-10. 2000 Ground-Water Quality Analyses of the Piedmont/Blue Ridge Aquifer System.

GWN-BR1B			
Well Name:	Young Harris New Well		
County:	Towns		
Date Sampled:	07/13/2000		
Nitrate/Nitrite	0.021	ppm	
pH	7.08	su	
conductivity	111	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-BR2A			
Well Name:	Notla Water Authority #3		
County:	Union		
Date Sampled:	07/13/2000		
Nitrate/Nitrite	0.96	ppm	
pH	4.83	su	
conductivity	34	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-BR4			
Well Name:	Morganton Old Well		
County:	Fannin		
Date Sampled:	09/14/2000		
Nitrate/Nitrite	1.6	ppm	
pH	5.44	su	
conductivity	81	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P1			
Well Name:	Luthersville New Well		
County:	Meriwether		
Date Sampled:	05/18/2000		
Nitrate/Nitrite	0.04	ppm	
pH	5.65	su	
conductivity	114	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	8.0	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	6.7	ppb	
Other			

GWN-P1			
Well Name:	Luthersville New Well		
County:	Meriwether		
Date Sampled:	07/19/2000		
Nitrate/Nitrite	--	ppm	
pH	5.47	su	
conductivity	86	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	5.5	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	6.0	ppb	
Other			

GWN-P1 [treated water]			
Well Name:	Luthersville New Well		
County:	Meriwether		
Date Sampled:	07/19/2000		
Nitrate/Nitrite	--	ppm	
pH	5.46	su	
conductivity	93	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	5.5	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds Trichloroethylene	3.8	ppb	
Other			

Table A-10 (Continued). 2000 Ground-Water Quality Analyses of the Piedmont/Blue Ridge Aquifer System.

GWN-P5			
Well Name:	Flowery Branch #1		
County:	Hall		
Date Sampled:	11/01/2000		
Nitrate/Nitrite	0.78	ppm	
pH	6.76	su	
conductivity	117	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P6A			
Well Name:	Shiloh #1		
County:	Harris		
Date Sampled:	04/06/2000		
Nitrate/Nitrite	0.022	ppm	
pH	7.21	su	
conductivity	127	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P7			
Well Name:	Hampton #6		
County:	Henry		
Date Sampled:	05/18/2000		
Nitrate/Nitrite	0.29	ppm	
pH	6.02	su	
conductivity	115	uS/cm	
Trihalomethanes: chloroform	0.59	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P8			
Well Name:	Wayne Farms #4		
County:	Jackson		
Date Sampled:	11/01/2000		
Nitrate/Nitrite	0.34	ppm	
pH	6.78	su	
conductivity	131	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P9A			
Well Name:	Gray #3		
County:	Jones		
Date Sampled:	03/09/2000		
Nitrate/Nitrite	0.06	ppm	
pH	6.46	su	
conductivity	359	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P10C			
Well Name:	Franklin Springs New #9		
County:	Franklin		
Date Sampled:	11/01/2000		
Nitrate/Nitrite	0.091	ppm	
pH	6.62	su	
conductivity	170	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other:			

Table A-10 (Continued). 2000 Ground-Water Quality Analyses of the Piedmont/Blue Ridge Aquifer System.

GWN-P11A			
Well Name:	Danielsville #2		
County:	Madison		
Date Sampled:	11/01/2000		
Nitrate/Nitrite	0.32	ppm	
pH	6.42	su	
conductivity	92	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other:	nd	ppb	

GWN-P12A			
Well Name:	Indian Spring		
County:	Butts		
Date Sampled:	03/16/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.25	su	
conductivity	223	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			
Fluoride	3.8	ppm	
Chloride	12	ppm	
Sulfate	65	ppm	

GWN-P13A			
Well Name:	Academy Spring/Covington		
County:	Newton		
Date Sampled:	03/16/2000		
Nitrate/Nitrite	0.74	ppm	
pH	6.07	su	
conductivity	53	uS/cm	
Trihalomethanes: chloroform	0.76	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P14			
Well Name:	Upson County/Sunset Village Well		
County:	Upson		
Date Sampled:	04/06/2000		
Nitrate/Nitrite	0.36	ppm	
pH	4.89	su	
conductivity	18	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P15A			
Well Name:	Bolton garden well		
County:	DeKalb		
Date Sampled:	03/16/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.14	su	
conductivity	163	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	0.78	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-P16C			
Well Name:	Mt. Airy #4		
County:	Habersham		
Date Sampled:	07/13/2000		
Nitrate/Nitrite	0.20	ppm	
pH	5.66	su	
conductivity	38	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds			
Vinyl chloride	1.1	ppb	
Other			

Table A-10 (Continued). 2000 Ground-Water Quality Analyses of the Piedmont/Blue Ridge Aquifer System.

GWN-P16C			
Well Name:	Mt. Airy #4		
County:	Habersham		
Date Sampled:	08/17/2000		
Nitrate/Nitrite	--		ppm
pH	5.57		su
conductivity	37		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-P17			
Well Name:	Oconee County/New Hillcrest Well		
County:	Oconee		
Date Sampled:	10/31/2000		
Nitrate/Nitrite	0.033		ppm
pH	7.23		su
conductivity	151		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-P18			
Well Name:	Dawsonville City Spring		
County:	Dawson		
Date Sampled:	09/14/2000		
Nitrate/Nitrite	1.4		ppm
pH	5.05		su
conductivity	49		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	0.51		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-P19			
Well Name:	Fayetteville #1		
County:	Fayette		
Date Sampled:	12/19/2000		
Nitrate/Nitrite	0.072		ppm
pH	7.15		su
conductivity	256		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-P20			
Well Name:	Suwanee #1		
County:	Gwinette		
Date Sampled:	12/20/2000		
Nitrate/Nitrite	0.35		ppm
pH	7.96		su
conductivity	158		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

Table A-11. 2000 Ground-Water Quality Analyses of the Valley and Ridge Aquifer System.

GWN-VR1			
Well Name:	Floyd County/Kingston Road Well		
County:	Floyd		
Date Sampled:	05/16/2000		
Nitrate/Nitrite	0.66	ppm	
pH	7.46	su	
conductivity	214	uS/cm	
Trihalomethanes:	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-VR2A			
Well Name:	LaFayette/Lower Big Spring		
County:	Walker		
Date Sampled:	05/17/2000		
Nitrate/Nitrite	1.8	ppm	
pH	7.45	su	
conductivity	252	uS/cm	
Trihalomethanes:	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-VR3			
Well Name:	Crawfish Spring/Chickamauga		
County:	Walker		
Date Sampled:	05/17/2000		
Nitrate/Nitrite	0.76	ppm	
pH	7.29	su	
conductivity	215	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-VR4			
Well Name:	Coats American #4		
County:	Walker		
Date Sampled:	05/17/2000		
Nitrate/Nitrite	nd	ppm	
pH	7.10	su	
conductivity	500	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-VR5			
Well Name:	Chattooga County #4		
County:	Chattooga		
Date Sampled:	05/17/2000		
Nitrate/Nitrite	2.6	ppm	
pH	6.93	su	
conductivity	342	uS/cm	
Trihalomethanes	nd	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds	nd	ppb	
Other			

GWN-VR6			
Well Name:	Chemical Products East Well		
County:	Bartow		
Date Sampled:	11/08/2000		
Nitrate/Nitrite	0.86	ppm	
pH	7.75	su	
conductivity	184	uS/cm	
Trihalomethanes: Chloroform	0.56	ppb	
Methyl tert-butyl ether	nd	ppb	
Benzene	nd	ppb	
Toluene	nd	ppb	
Ethylbenzene	nd	ppb	
Total Xylenes	nd	ppb	
Other Volatile Organic Compounds:			
1,1-Dichloroethylene	1.7	ppb	
Tetrachloroethylene	2.8	ppb	
Other			



Table A-11 (Continued). 2000 Ground-Water Quality Analyses of the Valley and Ridge Aquifer System.

GWN-VR7			
Well Name:	Adairsville/Lewis Spring		
County:	Bartow		
Date Sampled:	05/16/2000		
Nitrate/Nitrite	0.46		ppm
pH	7.61		su
conductivity	232		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-VR8			
Well Name:	Cedartown Spring		
County:	Polk		
Date Sampled:	05/16/2000		
Nitrate/Nitrite	0.68		ppm
pH	7.33		su
conductivity	240		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	1.2		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

GWN-VR9			
Well Name:	Polk County #2		
County:	Polk		
Date Sampled:	05/16/2000		
Nitrate/Nitrite	0.93		ppm
pH	7.18		su
conductivity	242		uS/cm
Trihalomethanes	nd		ppb
Methyl tert-butyl ether	nd		ppb
Benzene	nd		ppb
Toluene	nd		ppb
Ethylbenzene	nd		ppb
Total Xylenes	nd		ppb
Other Volatile Organic Compounds	nd		ppb
Other			

The Department of Natural Resources is an equal opportunity employer and offers all persons the opportunity to compete and participate in each area of DNR employment regardless of race, color, religion, national origin, age, handicap, or other non-merit factors.