



GEORGIA

DEPARTMENT OF NATURAL RESOURCES

ENVIRONMENTAL PROTECTION DIVISION

Canadian Wildfire Exceptional Event Demonstration for the 2024 Annual PM_{2.5} NAAQS at Columbus, GA in 2022-2024

Prepared by:
Air Protection Branch
Environmental Protection Division

September 19, 2025

TABLE OF CONTENTS

1. INTRODUCTION	1
2. NARRATIVE CONCEPTUAL MODEL	2
3. PUBLIC NOTIFICATION.....	3
4. CLEAR CAUSAL RELATIONSHIP AND SUPPORTING ANALYSES	5
5. NOT REASONABLY CONTROLLABLE OR PREVENTABLE	8
6. HUMAN ACTIVITY UNLIKELY TO RECUR AT A PARTICULAR LOCATION OR NATURAL EVENT	8
7. PUBLIC COMMENT PERIOD.....	9
APPENDIX A: ACTIVE WILDFIRES IN CANADA	8
APPENDIX B: HMS SMOKE AND ACTIVE FIRES.....	10
APPENDIX C: HYPLSIT BACK-TRAJECTORY MAPS	12
APPENDIX D: PM_{2.5} SURFACE CONCENTRATIONS IN THE SOUTHEAST	14
APPENDIX E: HOURLY PM_{2.5} TIME SERIES.....	16
APPENDIX F: UPPER AIR MAPS.....	18
APPENDIX G: PUBLIC COMMENTS	23
APPENDIX H: RESPONSES TO PUBLIC COMMENTS	23

1. Introduction

The current annual and 24-hour PM_{2.5} National Ambient Air Quality Standards (NAAQS) are 9.0 µg/m³ and 35 µg/m³, respectively. Federal Reference Method (FRM) monitors collect PM_{2.5} samples for 24 hours on filters while Federal Equivalent Method (FEM) monitors measure hourly PM_{2.5} concentrations continuously. For the purpose of this document, an “exceedance” is defined as a measured 24-hour PM_{2.5} concentration that is greater than the level (9.0 µg/m³) of the 2024 annual PM_{2.5} NAAQS. Please note that “exceedance” as defined in this document (based on an averaging time of 24-hours) is not an actual exceedance of the 2024 annual PM_{2.5} NAAQS since an exceedance of a NAAQS must be based on the averaging time for the NAAQS (in this case, annual) in addition to the level of the NAAQS (in this case, 9.0 µg/m³).

From 2019 through July 2020, PM_{2.5} data was collected at the Columbus-Cusseta site (AQS ID: 13-215-0011) with an FRM monitor on a one-in-three day schedule. This location was shut down at the end of July 2020, and then a new site was established in March 2021 at the Columbus-Baker site (AQS ID: 13-215-0012). The data from these two locations in Columbus (Cusseta and Baker) is linked in the U.S. Environmental Protection Agency (EPA) Air Quality System (AQS) for attainment purposes. At the Columbus-Baker location, one FRM monitor began collecting data in March 2021 on a one-in-three day sampling schedule. From 2021 through 2023, one FRM monitor operated at the Columbus-Baker site. In addition, an FEM monitor started collecting data in June 2023 and continued through 2024, with a NAAQS exclusion on the data. The Columbus, GA-AL Metropolitan Statistical Area (MSA) is in attainment of the 2012 PM_{2.5} NAAQS.

This document discusses 10 different days from 2022-2024 that qualify for exceptional event demonstrations (Table 1) for the Columbus-Baker air monitoring site (AQS ID: 13-215-0012) located in Columbus (Muscogee County) in the state of Georgia. Of these events, 2 were due to Canadian wildfires. These two exceedances resulted from the transport of wildfire smoke that originated in Canada. Therefore, they qualify for removal under the Exceptional Events Rule (EER). Design values (DVs) of the Columbus-Baker site with and without EPA concurrence are shown in Table 2. DVs are calculated using 24-hour PM_{2.5} measurements from 2022-2024. For each year, these measurements are first averaged into quarterly values, then to a yearly value. The average of these yearly values is reported as the DV. Inclusion of these events produces a DV of 9.5 µg/m³, which violates the new 2024 PM_{2.5} annual NAAQS; however, exclusion reduces the DV to 9.0 µg/m³.

On July 30, 2025, the Georgia Environmental Protection Division (EPD) submitted an Initial Notification for these events to EPA. The request indicated that two events identified in Table 1 were impacted by smoke from wildfires and requested review of the events under the case-by-case provision at 40 CFR 50.14(a)(1)(i)(F). This demonstration will focus on the two Canadian wildfire events, while a separate demonstration will focus on the eight prescribed fire events. Georgia EPD formally requests that EPA concur with the exclusion of these wildfire events.

Table 1. Exceedances observed at the Columbus-Baker site in Columbus, GA in 2022-2024 that qualify for removal under the EER.

#	Date	24-hour PM _{2.5} (µg/m ³)	Tier	Cause of Exceedance
1	02/01/22	22.1	1	Prescribed fires
2	02/10/22	32.0	1	Prescribed fires
3	03/03/22	55.1	1	Prescribed fires
4	03/01/23	21.1	1	Prescribed fires
5	03/07/23	25.0	1	Prescribed fires
6	06/29/23	20.8	1	Canadian wildfires
7	07/17/23	25.3	1	Canadian wildfires
8	02/03/24	34.3	1	Prescribed fires
9	02/21/24	28.5	1	Prescribed fires
10	03/13/24	35.0	1	Prescribed fires

Table 2. Design values at the Columbus-Baker site for the 2024 annual PM_{2.5} NAAQS.

Monitor Site (AQS ID)	2022-2024 DV without EPA Concurrence (µg/m ³)	2022-2024 DV with EPA Concurrence (µg/m ³)
Columbus-Baker (13-215-0012)	9.6	9.0

The EPA has outlined requirements for demonstrations of wildfire events in the 2016 document *Guidance on the Preparation of Exceptional Events Demonstrations for Wildfire Events that May Influence Ozone Concentrations* and, pertinent to this demonstration, the 2024 supplementary document *PM_{2.5} Wildland Fire Exceptional Events Tiering Document*. This demonstration will describe how the proposed wildfire events meet the requirements of the EER as described in regulation and the guidance documents, as applicable.

2. Narrative Conceptual Model

The EER requires that demonstrations include a narrative conceptual model describing the event. This section describes the 2023 wildfires that impacted air quality in Columbus, GA. Estimates from the National Oceanic and Atmospheric Administration (NOAA) Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT) model are used to describe the transport of wildfire smoke to the area and around the state which ultimately led to enhancements of PM_{2.5} concentrations that exceeded the NAAQS level.

Canadian wildfires during the 2023 wildfire season were well documented and impacted much of the geography of the United States. This season started ahead of the typical Canadian wildfire season, lasting from mid-April to late October (seasons are usually from May – September)¹. Temperatures and land aridity across Canada were unusually high and resulted in the burning of a record-breaking amount of land area (≥156,000 km²)². The land area burned during this season far exceeded the average³ of 21,000 km², with the most active burns situated in the eastern province of Quebec in June and July.

¹ <https://doi.org/10.1038/s41467-024-51154-7>

² <https://doi.org/10.1007/s00376-023-3241-0>

³ <https://cwfis.cfs.nrcan.gc.ca/ha/nfdb>

Figures in Appendix A are provided to show active Canadian wildfires on the days of the exceedances and for three days beforehand via the Natural Resources Canada Interactive Map⁴. For Figures A1 and A2, when the exceedances were recorded at the relevant site, Canadian wildfires were on-going across the country, the majority of which had each consumed >1,000 hectares. These fires were similarly as intense up to three days prior to the recorded exceptional events.

Shown in Section 4, these wildfires resulted in the United States being blanketed in smoke and impacted PM_{2.5} surface level concentrations across the country. Pertinent to this demonstration, concentrations were impacted across the southeast of the country, often simultaneously and in conjunction with the arrival of air masses either from Canada or circulated from smoke-laden areas within the United States. Air mass back-trajectories from NOAA's HYSPLIT model indicate that the plumes responsible for the summertime events were emitted from fires in the Canadian provinces of British Columbia, Alberta, Saskatchewan, and Quebec. They then traveled along multiple routes either originating in Canada or by circulating smoke from Canadian wildfires from other (western and/or midwestern) regions of the United States.

This conceptual model describes how emissions from wildfires in Canada and environmental conditions contributed to the wildfire events dated in Table 1. Smoke emissions transported to the Columbus-Baker site enhanced observed PM_{2.5} concentrations and caused an exceedance. Georgia EPD requests EPA's concurrence on June 29 and July 17, 2023, for exclusion from regulatory decision making, specifically state attainment determinations.

3. Public Notification

As described in 40 CFR 51.930(a), states requesting to exclude data due to exceptional events must take appropriate and reasonable actions to protect public health from exceedances or violations of the NAAQS. These include providing for, at a minimum, prompt public notification whenever concentrations are expected to exceed a NAAQS, public education on actions individuals may take to reduce exposures to unhealthy air quality during events, and implementation of appropriate measures to protect public health from event-caused exceedances or violations of the NAAQS.

With respect to public notification and public education, the Georgia Forestry Commission (GFC) has a public website⁵ with an interactive wildfire and burn permit map that contains the current Air Quality Index at all sites in Georgia with the option to add the following layers: (1) burn restrictions, (2) daily burn permits, (3) PM_{2.5}, (4) NOAA Hazard Mapping System (HMS) smoke plumes, (5) wind vectors, and (6) smoke forecast. The public can zoom in to see if smoke may impact their location. The Georgia EPD website⁶ has a link to the GFC interactive burn permit map. Also, the Georgia EPD website has a link to EPA's AirNow Fire and Smoke Map⁷,

⁴ <https://cwfis.cfs.nrcan.gc.ca/interactive-map>

⁵ <https://georgiafc.firesponse.com/public/>

⁶ <https://epd.georgia.gov/air-protection-branch/open-burning-rules-georgia>

⁷ <https://fire.airnow.gov/>

EPA's AirNow When Smoke is in the Air⁸, EPA's AirNow Prepare for Fire Season⁹, and the EPA's Smoke-Ready Toolbox for Wildfires¹⁰. These websites identify several protective measures that individuals should take to reduce smoke exposure as needed, including limiting outdoor activities, avoiding strenuous outdoor activity and remaining indoors, and considering temporarily relocating or closing all doors and windows during smoke events. In addition, the Georgia EPD Ambient Air Monitoring Program website¹¹ provides near real-time ambient air concentrations of multiple criteria pollutants (O₃, PM_{2.5}, SO₂, NO₂, and CO) across the state.

Georgia EPD, in conjunction with the Georgia Institute of Technology, provides a daily forecast email that includes messaging concerning health advisories and smog alerts to Georgia Commute Options and the Atlanta Regional Commission for distribution. Each forecast email includes a reminder stating: "If you see or smell smoke and have respiratory concerns you may need to move indoors, close windows and doors. You can see the current air quality at <https://airgeorgia.org> or <https://airnow.gov> to help you determine when to continue outdoor activities."

Outside Georgia EPD and GFC, there was additional notification provided by various news outlets to the public about the potential for elevated air quality impacts from Canadian wildfires. Some examples include:

- <https://www.wsbtv.com/news/local/atlanta/expect-flight-delays-into-atlanta-thanks-canada-wildfire-smoke/RX2WTAWHTBGEVQAMWFRLYCEJU/> (North Georgia/Metro Atlanta, 6/7/2023)
- <https://www.13wmaz.com/article/weather/smoke-from-wildfire-to-central-georgia/93-87f7b552-98fd-4b86-8f99-94140c76d38d> (Macon/Central Georgia, 6/7/2023)
- https://www.youtube.com/watch?v=sZkxex_jpQw (Central Georgia, 6/7/2023)
- <https://www.wsbradio.com/weather/wildfire-smoke-drifting-south-canada-into-north-georgia/OKNENTCA6JGGJPYQ76VCO6LF5Q/> (Metro Atlanta/Birmingham, Alabama, 6/7/2023)
- <https://www.gpb.org/news/the-picture-show/2023/06/07/photos-extreme-canadian-wildfire-smoke-shrouds-parts-of-us> (North and South Carolina, 6/7/2023)
- <https://www.atlantanevnewsfirst.com/2023/06/08/air-quality-georgia-will-smoke-wildfires-reach-us/> (Greater Metro Atlanta, 6/8/2023)
- <https://patch.com/georgia/atlanta/code-orange-alert-what-know-air-quality-ga> (Atlanta, 6/8/2023)
- <https://www.walb.com/2023/06/08/canadian-wildfire-smoke-arrives-south-ga-effects-are-minimal/> (Albany/South Georgia, 6/8/2023)
- <https://abcnews.go.com/US/canadian-wildfire-dangers-prompt-proactive-mitigation-government-experts/story?id=100478859> (Predicts plume will migrate to Georgia, 7/1/2023)
- <https://www.11alive.com/article/weather/stormtracker/wildfire-smoke-north-georgia/85-b4670fee-4608-4f99-9904-bbdbc924375> (North Georgia, 7/17/2023)

⁸ <https://www.airnow.gov/wildfires/when-smoke-is-in-the-air/>

⁹ <https://www.airnow.gov/sites/default/files/2020-10/prepare-for-fire-season.pdf>

¹⁰ <https://www.epa.gov/air-research/smoke-ready-toolbox-wildfires>

¹¹ <https://airgeorgia.org/>

- <https://foxchattanooga.com/weather/stormtrack-9-blog/canadian-wildfire-smoke-impacts-tennessee-georgia-air-quality-once-again> (Chattanooga, 7/17/2023)
- <https://www.nytimes.com/2023/07/18/us/smoke-wildfires-nc-georgia.html> (Georgia, 7/18/2023)
- <https://www.fox5atlanta.com/news/code-orange-alert-metro-atlanta-canadian-wildfire-smoke> (Georgia, 7/18/2023)
- <https://www.atlantaneewsfirst.com/2023/07/18/canadian-wildfires-bring-poor-air-quality-north-georgia/> (Metro Atlanta/North Georgia, 7/18/2023)
- <https://www.cnn.com/2023/07/17/weather/canada-wildfires-shatter-burning-records/index.html> (North Georgia, 7/18/2023)
- <https://www.iqair.com/us/newsroom/atlanta-air-quality-alert> (Atlanta 7/18/2023)
- <https://www.houstonchronicle.com/news/houston-texas/environment/article/houston-pollution-canadian-wildfire-saharan-dust-18206844.php> (Atlanta/Georgia, 7/18/2023)
- <https://www.wsfa.com/2023/07/18/details-behind-canadian-wildfire-smoke-alabamas-sky/> (Montgomery/Alabama, 7/18/2023)
- <https://news.gatech.edu/news/2023/07/19/canadian-wildfire-smoke-affects-atlanta-2> (Atlanta, 7/19/2023)
- <https://www.gpb.org/news/2023/07/26/macon-had-georgias-worst-air-quality-last-week-thanks-canadian-wildfires-heat-dome> (Macon, 7/26/2023)
- <https://www.savannahnow.com/story/weather/2023/10/03/savannah-ga-air-quality-canadian-wildfire-smoke-moves-down-u-s-coast/71045920007/> (Savannah, 10/3/2023)
- <https://www.gpb.org/news/2023/10/04/have-you-noticed-haze-in-the-air-heres-why-georgia-dealing-smoke> (South/Middle Georgia, 10/4/2023)

4. Clear Causal Relationship and Supporting Analyses

This section addresses the EER requirements at 40 CFR 50.14(c)(3)(iv)(B) by showing that the events affected air quality in such a way that there exists a clear, causal relationship between the specific events and the monitored exceedance, and at 40 CFR 50.14(c)(3)(iv)(C) by providing analyses comparing the claimed event-influenced concentrations to concentrations at the same monitoring site at other times. The *Guidance on the Preparation of Exceptional Events Demonstrations for Wildfire Events that May Influence Ozone Concentrations and PM_{2.5} Wildland Fire Exceptional Events Tiering Document* outlines the expected components of a clear causal relationship portion of a demonstration. These include a comparison of the event-related concentration to historical concentrations, evidence that the emissions from wildfires were transported to the site, and evidence that the prescribed fire emissions affected the monitor.

Figures B1 and B2 (Appendix B) show smoke from the NOAA Hazard Mapping System (HMS), plotted via the AirNow Navigator¹². Active fires and smoke are shown for the day the event was registered as well as up to three days beforehand. During the exceptional events, smoke pervaded the air throughout much of the eastern half of the United States, if not the entire country.

The historical data analysis section of this demonstration focuses on 2020-2024 PM_{2.5} FRM data at the Columbus-Baker site. Table 3 contains a comparison of exceptional event concentrations

¹² <https://airnowtech.org/navigator/>

to historic 2020-2024 concentrations for the site. Generally, the exceptional event concentrations are at least double the 5-year annual average, quarterly average, and monthly average, and in some cases can be up to three times higher.

Table 3. Comparison of exceptional event concentrations to historic 2020-2024 concentrations at the Columbus-Baker site.

EE Date	EE Concentration ($\mu\text{g}/\text{m}^3$)	5-Year Annual Average ($\mu\text{g}/\text{m}^3$)	5-Year Quarterly Average ($\mu\text{g}/\text{m}^3$)	5-Year Monthly Average ($\mu\text{g}/\text{m}^3$)	Ratio EE to 5-Year Annual Average	Ratio EE to 5-Year Quarterly Average	Ratio EE to 5-Year Monthly Average
06/29/23	20.8	9.2	8.75	9.56	2.3	2.4	2.2
07/17/23	25.3	9.2	8.37	8.55	2.8	3.0	3.0

Figure 1 plots the 24-hour $\text{PM}_{2.5}$ concentrations for 2020-2024. Exceedances caused by wild or prescribed fires are delineated by marker shape. The selected exceptional events are above the Tier 1 threshold of $17.7 \mu\text{g}/\text{m}^3$. Tier 1 events are 1.5 times greater than the highest 98th percentile of data over the last 5 years, per the EPA’s Tiering Tool.




Figure 1. 24-hour $\text{PM}_{2.5}$ concentrations for 2020-2024 observed at the Columbus-Baker site. Exceedances not related to fires are demarcated by color with Tier 1 in red, Tier 2 in yellow, and Tier 3 in blue. Canadian wildfires (green squares) and prescribed burns (green circles) are additionally differentiated.

Maps from the AirNow Navigator are provided in Appendix C for each exceedance event in Table 3. The maps include NOAA HMS satellite detected fires, HMS smoke plumes, 24-hour rolling average PM_{2.5} concentrations (including the time shown in the figure panel and the 23 hours following it) across the United States, and HYSPLIT back-trajectories. These trajectories originate at the Columbus-Baker site's geographic location and extend 72 hours back in time. Three different starting times were modeled with HYSPLIT for each day: (1) midnight at the start of the exceedance day, (2) noon of the exceedance day, and (3) midnight at the end of the exceedance day. Each trajectory starts at a different elevation above ground level (100 m, 1500 m, and 3000 m). The 1500 m and 3000 m tails are used to estimate the trajectories of smoke transported over a long range. The 100 m tail, nearer to the surface, is used to show local transport. Additionally, these values are chosen to estimate vertical transport near-surface and up to several hundred meters above the planetary boundary layer. Figures in Appendix D are provided to show ground level, daily PM_{2.5} concentrations, and air quality indices (AQIs) in the southeast of the United States. Figures in Appendix E show hourly PM_{2.5} time series for both the event day and the day before. Figures in Appendix F show upper air maps from the Storm Prediction Center¹³ for the event day and three days prior. Maps are displayed at pressures of either 850 millibar (mb), equivalent to 1170-1590 m above mean sea level (MSL), or 700 mb, equivalent to 2350-3150 MSL¹⁴, at 00 Coordinated Universal Time (UTC) or 12 UTC for each day. These pressure values are chosen to correspond with the 1500 m and 3000 m heights of the HYSPLIT trajectories. A specific pressure value is determined on a case-by-case basis depending on how clearly the corresponding upper air maps explain the sequence of events that led to the relevant exceedance.

June 29, 2023

Fires in Canada had been on-going for months at the time the exceedance was measured, and their emissions were likely mixed throughout the air column. From Figure E1, PM_{2.5} concentrations were elevated above the 9.0 µg/m³ standard throughout the event day and the day prior. Shown in Figure C1, the 1500 m HYSPLIT tail, released on midnight of the exceedance day, traveled from near a large plume in the northeast U.S. emitted by the Quebec fires. The 1500 m tails released at later times on the event day follow similar spatial trajectories, indicating the Quebec smoke plume was transported to the site. Wind barbs in Figures F1 and F2 corroborate the back-trajectories as they show a counterclockwise vortex centered over the northeast/Great Lakes region on June 26 and 27 that extended to the southeast. Smoke plumes were initially circulated in this vortex and then advected through the Midwest and down to the southeast (Figures F3-F5). This led to the daily PM_{2.5} concentration at the Columbus-Baker site on June 29, 2023, increasing to 20.8 µg/m³. Figure D1 shows that similar enhancements occurred synchronously at other sites across the southeast, which follows from the large blanket of smoke over this region (Figure B1).

July 17, 2023

Fires in Canada had been on-going for months at the time the exceedance was measured, and their emissions were likely mixed throughout the air column. From Figure E2, PM_{2.5} concentrations were elevated above the 9.0 µg/m³ standard starting on 10:00 PM EST on July 16 and remained elevated throughout the event day. Shown in Figure C2, back-trajectories indicate

¹³ <https://www.spc.noaa.gov/obswx/maps/>

¹⁴ <https://www.noaa.gov/jetstream/upper-air-charts>

that the smoke plume traveled through the Midwest of the United States. As a result, the plume detected at the site is most likely a mixture of emissions from fires in the provinces of British Columbia, Alberta, and Saskatchewan. The back-trajectories converge in space approximately 24 hours before descending to near-surface level where observed, the daily PM_{2.5} concentration at the Columbus-Baker site on July 17, 2023, increased to 25.3 µg/m³. Figure D2 shows that similar enhancements occurred synchronously at other sites across the southeast, which follows from the large blanket of smoke over this region (Figure B2).

The comparisons and analyses provided in this demonstration support Georgia EPD's position that the fire event affected air quality in such a way that there exists a clear causal relationship between the specific event and the monitored exceedance for the dates described in Table 1 and thus satisfies the clear causal relationship criterion.

5. Not reasonably Controllable or Preventable

This section satisfies the EER requirements at 40 CFR 50.14(c)(3)(iv)(A), CFR 50.1(j), 40 CFR 50.14(c)(3)(iv)(D), and 40 CFR 50.14(b)(4): The event was caused by a natural event; an exceptional event is one that is not reasonably controllable or preventable. Stated in section 40 CFR 50.14 (a)(8)(vii), the Administrator shall not require a State to provide case-specific justification to support the not reasonably controllable or preventable criterion for emissions-generating activity that occurs outside of the State's jurisdictional boundaries within which the concentration at issue was monitored.

6. Human Activity Unlikely to Recur at a Particular Location or Natural Event

This section satisfies the EER requirement at 40 CFR 50.14(c)(3)(iv)(E): A demonstration that the event was a human activity that is unlikely to recur at a particular location or was a natural event. The definition of wildfire in the EER is: "...any fire started by an unplanned ignition caused by lightning; accidental, human causes actions, or a prescribed fire that has developed into a wildfire. A wildfire that predominately occurs on wildland is a natural event." As stated in sections 2 and 4, the origin and evolution of the wildfires described in this demonstration occurred in Canada.

Based on the documentation provided in Sections 2 and 4 of this demonstration, these events qualify as natural events as they spread uncontrolled through remote, natural (i.e., non-agricultural or silvicultural) lands. The National Aeronautics and Space Administration noted that many of the Canadian fires were ignited by summer lightning storms and largely burned in deeply wooded areas¹⁵. EPA generally considers the emissions of PM_{2.5} from wildfires to meet the regulatory definition of a natural event, defined as one "in which human activity plays little or no direct causal role" (40 CFR 50.1(k)). As Georgia EPD has shown that the demonstrated exceedances resulted from natural events, they should be considered for treatment as exceptional events.

¹⁵ <https://earthobservatory.nasa.gov/images/151985/tracking-canadas-extreme-2023-fire-season>

7. Public Comment Period

Georgia EPD held a public comment period beginning August 4, 2025, to receive public input regarding the Exceptional Event Demonstration. Notification of the public comment period was posted on the Georgia EPD website and emailed to interested stakeholders. Public comments received are included in Appendix G of this document, along with Georgia EPD's responses to these comments in Appendix H.

Appendix A: Active Wildfires in Canada







Figure A1. Active wildfires in Canada on June 26-29, 2023.







Figure A2. Active wildfires in Canada on July 14-17, 2023.

Appendix B: HMS Smoke and Active Fires




Figure B1. Map from the AirNow Navigator showing active fires (red triangles) and smoke (grey polygons) on June 26-29, 2023, over North America.







Figure B2. Map from the AirNow Navigator showing active fires (red triangles) and smoke (grey polygons) on July 14-17, 2023, over North America.

Appendix C: HYPLSIT Back-Trajectory Maps




Figure C1. Map of HMS smoke plumes (grey polygons) and fires (red triangles), 24-hour rolling average PM_{2.5} concentrations (circles), and HYSPLIT back-trajectories of release heights at 100 m (green lines), 1500 m (blue lines), and 3000 m (red lines) for 0 AM EST on June 29, 2023 (top), 12 PM EST on June 29, 2023 (middle), and 0 AM EST on June 30, 2023 (bottom).





Figure C2. Map of HMS smoke plumes (grey polygons) and fires (red triangles), 24-hour rolling average PM_{2.5} concentrations (circles), and HYSPLIT back-trajectories of release heights at 100 m (green lines), 1500 m (blue lines), and 3000 m (red lines) for 0 AM EST on July 17, 2023 (top), 12 PM EST on July 17, 2023 (middle), and 0 AM EST on July 18, 2023 (bottom).

Appendix D: PM_{2.5} Surface Concentrations in the Southeast June 29, 2023 PM_{2.5} Exceedance Report




AQI category - 24-hr average PM_{2.5}

- Good (0-9 ug/m³)
- Moderate (9.1-35.4 ug/m³)
- Unhealthy for sensitive (35.5-55.4 ug/m³)
- Unhealthy (55.5-150.4 ug/m³)
- Very unhealthy (150.4-250.4 ug/m³)
- Hazardous (>250.4 ug/m³)

Figure D1. Surface level, daily PM_{2.5} concentrations on June 29, 2023, across the southeast. The Columbus-Baker site is represented by a star. Numerous sites measured concentrations that exceeded the level of annual PM_{2.5} NAAQS.

July 17, 2023 PM_{2.5} Exceedance Report



AQI category - 24-hr average PM_{2.5}

- Good (0-9 ug/m³)
- Moderate (9.1-35.4 ug/m³)
- Unhealthy for sensitive (35.5-55.4 ug/m³)
- Unhealthy (55.5-150.4 ug/m³)
- Very unhealthy (150.4-250.4 ug/m³)
- Hazardous (>250.4 ug/m³)

Figure D2. Surface level, daily PM_{2.5} concentrations on July 17, 2023, across the southeast. The Columbus-Baker site is represented by a star. Numerous sites measured concentrations that exceeded the level of annual PM_{2.5} NAAQS.

Appendix E: Hourly PM_{2.5} Time Series




Figure E1. Hourly PM_{2.5} concentrations at the Columbus-Baker site on June 28 and 29, 2023. The solid lines with dots show observations, and the dashed lines show AQI tiers adjusted for the 2024 PM_{2.5} NAAQS.




Figure E2. PM_{2.5} concentrations at the Columbus-Baker site on July 16 and 17, 2023. The solid lines with dots show observations, and the dashed lines show AQI tiers adjusted for the 2024 PM_{2.5} NAAQS.

Appendix F: Upper Air Maps




Figure F1. Storm Prediction Center upper air maps for June 26, 2023, at 00 UTC (top) and at 12 UTC (bottom). Maps are generated at a pressure of 850 mb (altitude of 1170-1590 MSL). Wind barbs (degrees from north, knots) are plotted in blue, isotherms (degrees Celsius (°C)) in red, isodrosotherms (°C) in green contours, and isoheights (MSL) in dark grey.




Figure F2. Storm Prediction Center upper air maps for June 27, 2023, at 00 UTC (top) and at 12 UTC (bottom). Maps are generated at a pressure of 850 mb (altitude of 1170-1590 MSL). Wind barbs (degrees from north, knots) are plotted in blue, isotherms ($^{\circ}\text{C}$) in red, isodrosotherms ($^{\circ}\text{C}$) in green, and isoheights (MSL) in dark grey.




Figure F3. Storm Prediction Center upper air maps for June 28, 2023, at 00 UTC (top) and at 12 UTC (bottom). Maps are generated at a pressure of 850 mb (altitude of 1170-1590 MSL). Wind barbs (degrees from north, knots) are plotted in blue, isotherms (°C) in red, isodrosotherms (°C) in green, and isoheights (MSL) in dark grey.




Figure F4. Storm Prediction Center upper air maps for June 29, 2023, at 00 UTC (top) and at 12 UTC (bottom). Maps are generated at a pressure of 850 mb (altitude of 1170-1590 MSL). Wind barbs (degrees from north, knots) are plotted in blue, isotherms ($^{\circ}\text{C}$) in red, isodrosotherms ($^{\circ}\text{C}$) in green, and isoheights (MSL) in dark grey.




Figure F5. Storm Prediction Center upper air maps for June 30, 2023, at 00 UTC (top) and at 12 UTC (bottom). Maps are generated at a pressure of 850 mb (altitude of 1170–1590 MSL). Wind barbs (degrees from north, knots) are plotted in blue, isotherms (°C) in red, isodrosotherms (°C) in green, and isoheights (MSL) in dark grey.

Appendix G: Public Comments

Appendix H: Responses to Public Comments