Drought Indicators Report

Georgia Environmental Protection Division

January 2023

Background

Pursuant to the Rules for Drought Management, <u>Section 391-3-3-.04</u> Drought <u>Indicators and Triggers</u>, the Director of EPD monitors climatic indicators and water supply conditions to assess drought occurrence and severity, and its impact upon the ability of public water systems to provide adequate supplies of water. These indicators and conditions December include, but not be limited, to the following:

- U.S. Drought Monitor;
- Precipitation;
- Streamflow;
- Groundwater;
- Reservoir levels;
- Short term climate predictions;
- Soil moisture; and
- Water supply conditions.

Background

- The Rules require EPD to report on current climatic indicators at least semi-annually or monthly when any part of the state has experienced at least two consecutive months of severe drought.
- This reports compare current conditions to historical levels (and/or reservoir rule curves) for each of the following indicators:
 - Precipitation during the prior 3, 6, and 12 months;
 - Streamflow at the select United States Geological Survey gages;
 - Groundwater levels at select United States Geological Survey monitoring wells; and
 - Reservoir levels at Allatoona Lake, Lake Hartwell, Clarks Hill Lake, and Lake Lanier.
- The following sections of this presentation provide the data and information sources analyzed by EPD in developing this drought indicators report for conditions as of January 10, 2023.

Drought Indicator Analysis Summary (slide 1 of 2)


- U.S. Drought Monitor Severe drought (D2) exists in Seminole, Early, Miller, Decatur, Grady, Thomas, Brooks and Lowndes Counties. Moderate drought (D1) exists in Lower Flint Basin, Suwannee Basin and Coastal area. Abnormally Dry (D0, the least intense level) exists in most areas below Fall Line.
- Precipitation Three-month precipitation is below normal in most areas below Fall Line and some areas in North Georgia. Six-month precipitation is slightly below normal in Echols and Ware Counties and Satilla Basin. Twelve-month precipitation is below normal in most areas below Fall Line and Upper Savannah Basin.
- Soil Moisture Soil moisture conditions are below normal in Counties along state line with Florida.

Drought Indicator Analysis Summary (slide 2 of 2)

- Streamflow Stream flows at majority of selected USGS gages (19 out of 34) are near or above normal. 14 gages are between the lowest 20th percentile and median. One gage (in Ocmulgee Basin) is between the lowest 20th and 10th percentiles.
- Groundwater Level Groundwater levels are above or near normal in four wells (4 out of 17). Ten well levels are between the lowest 20th percentile and median (10 out of 17). Three well levels are between the lowest 20th and 10th percentiles (two in Flint Basin and one in Suwannee Basin).
- Reservoir Levels At the end of December 2022, all federal reservoirs in ACT Basin and Savannah Basin are at levels above or near their respective top of conservation (normal) pools. Federal reservoirs in ACF Basin are in zone 1. ACF composite storage is in zone 1.
- Short-term Climate Prediction National Climatic Prediction Center projects above normal temperature statewide and below normal precipitation in most areas in January – March 2023. U.S. Drought Outlook predicts drought persists in Coastal area, drought remains but improves in Southwest corner, and drought removal likely in other areas in January – March 2023.
- Water Supplies No issues with water availability to water supply providers were reported.

US Drought Monitor


Data Source: http://droughtmonitor.unl.edu/



U.S. Drought Monitor Georgia

January 3, 2023 (Released Thursday, Jan. 5, 2023) Valid 7 a.m. EST

Drought Conditions (Percent Area)




	None	D0-D4	D1-D4	D2-D4	D3-D4	D4
Current	46.36	53.64	28.04	4.81	0.00	0.00
Last Week 12-27-2022	46.36	53.64	27.82	4.81	0.00	0.00
3 Month s Ago 10-04-2022	70.82	29.18	2.48	0.00	0.00	0.00
Start of Calendar Year 01-03-2023	46.36	53.64	28.04	4.81	0.00	0.00
Start of Water Year 09-27-2022	76.20	23.80	0.00	0.00	0.00	0.00
One Year Ago 01-04-2022	97.01	2.99	0.00	0.00	0.00	0.00

Intensity:







D3 Extreme Drought

D1 Moderate Drought

D4 Exceptional Drought

The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. For more information on the Drought Monitor, go to https://droughtmonitor.unl.edu/About.aspx

Author:

Brad Pugh CPC/NOAA




droughtmonitor.unl.edu


3, 6, and 12 Month Precipitation Anomaly

Data Source: https://www.ncdc.noaa.gov/cag/county/mapping/


3 Month Precipitation Anomaly



6 Month Precipitation Anomaly




12 Month Precipitation Anomaly



Soil Moisture Conditions

Data Source: http://www.hydro.ucla.edu/SurfaceWaterGroup/forecast/moni tor/curr/conus.mexico/east.vic.sm_qnt.gif




Current (Daily updated) percentiles for soil moisture (SWE) with respect to the climatological period (1916-2004).

Streamflow Conditions

Data Source: USGS

Streamflow Monitoring

- As shown on the following slide, EPD Monitors 34 USGS stream gages in 13 of the State's major river basins to assess drought conditions.
- These gages were selected because each has:
 - Long-term and relatively complete records for recent decades; and
 - Relatively low consumptive water use implications and streamflows are not heavily influenced by dams.
- Note: Hydrologic conditions of major rivers with streamflows that are heavily influenced by dams can be assessed by reviewing status of major storage reservoirs



USGS Stream Gages Monitored by EPD to Assess Drought Conditions

GAGE#	BASIN	GAGE NAME
1	TENNESSEE	LOOKOUT CREEK NEAR NEW ENGLAND
2	TENNESSEE	NOTTELY RIVER NEAR BLAIRSVILLE
3	COOSA	CHATTOOGA RIVER AT SUMMERVILLE
4	COOSA	TALKING ROCK CREEK NEAR HINTON
5	COOSA	ETOWAH RIVER AT CANTON
6	CHATTAHOOCHEE	CHATTAHOOCHEE RIVER AT CORNELIA
7	CHATTAHOOCHEE	CHESTATEE RIVER NEAR DAHLONEGA
8	CHATTAHOOCHEE	NEW RIVER AT GA 100 NEAR CORINTH
9	CHATTAHOOCHEE	UPATOI CREEK AT COLUMBUS
10	FLINT	FLINT RIVER AT GA26 NEAR MONTEZUMA
11	FLINT	FLINT RIVER AT ALBANY
12	FLINT	ICHAWAYNOCHAWAY CREEK AT MILFORD
13	FLINT	SPRING CREEK NEAR IRON CITY
14	OCMULGEE	ALCOVY RIVER ABOVE COVINGTON
15	OCMULGEE	OCMULGEE RIVER AT MACON
16	OCMULGEE	TOBESOFKEE CREEK NEAR MACON
17	OCMULGEE	TUCSAWHATCHEE CREEK NEAR
		HAWKINSVILLE
18	OCONEE	MIDDLE OCONEE RIVER NEAR ATHENS
19	OCONEE	LITTLE RIVER NEAR EATONTON
20	OCONEE	OCONEE RIVER AT DUBLIN
21	ALTAMAHA	OHOOPEE RIVER NEAR REIDSVILLE
22	SAVANNAH	CHATTOOGA RIVER NEAR CLAYTON
23	SAVANNAH	BROAD RIVER NEAR BELL
24	SAVANNAH	BEAVERDAM CREEK NEAR SARDIS
25	SAVANNAH	BRIER CREEK AT MILLHAVEN
26	OGEECHEE	CANOOCHEE RIVER NEAR CLAXTON
27	OGEECHEE	OGEECHEE RIVER NEAR EDEN
28	OCHLOCKONEE	OCHLOCKONEE RIVER NEAR THOMASVILLE
29	SUWANEE	WITHLACOOCHEE RIVER NEAR PINETTA FL
30	SUWANEE	ALAPAHA RIVER AT STATENVILLE
31	SUWANEE	SUWANNEE RIVER AT US 441, AT FARGO
32	SATILLA	SATILLA RIVER NEAR WAYCROSS
33	SATILLA	LITTLE SATILLA RIVER NEAR OFFERMAN
34	ST MARY	ST MARYS RIVER NEAR MACCLENNY FL

Streamflow Graphs

- For each of the 34 gages, EPD has prepared a graph that shows monthly average streamflow from January 2022 through December 2022;
- To help put these streamflow conditions into perspective, for comparison purposes, each graph also shows:
 - Monthly average streamflows for the years 2007 and 2011 when streamflows were at or near recorded low levels across much of the state; and
 - A statistical composite of historical conditions showing the "driest" 50, 20, 10, and 5 percent of all recorded monthly average stream flows at the same gage.


How to Read the Streamflow Graphs Example #1: Etowah River at Canton


The streamflow graph for Gage #5, <u>USGS Etowah River gage at Canton</u> shows :


- Average stream flow in December 2022 was 1642 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2022 about 80% of the time; 20% of the time in December it has been higher.
- Average stream flow in December 2011 was 790 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2011 about 30~40% of the time; 60~70% of the time in December it has been higher.
- Average stream flow in December 2007 was 342 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2007 about 0.1 % of the time; 99.9% of the time in December it has been higher.


How to Read the Streamflow Graphs <u>Example #2:</u> Flint River at Albany


- The streamflow graph for Gage #11, <u>USGS Flint River gage at Albany</u> shows:
- Average stream flow in December 2022 was 4445 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2022 about 51% of the time; about 49% of the time in December it has been higher.
- Average stream flow in December 2011 was 2100 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2011 about 2~5% of the time; about 95~98% of the time in December it has been higher.
- Average stream flow in December 2007 was 2463 cfs. The statistical composite of all historical data for this gage shows that average streamflow in December has historically been lower than December 2007 about 10% of the time; about 90% of the time in December it has been higher.


































































































Groundwater Levels

Data Source: USGS

Rationale for Choosing USGS Monitoring Wells

EPD monitors 17 groundwater USGS monitoring wells shown on the following slide to assess drought conditions. These wells were selected for monitoring because they have:

- Long-term monitoring records consisting of three decades or more of data; and
- Real-time monitoring that represents the most up-to-date conditions.

USGS Wells Monitored

Chattahoochee Basin

1.16MM03

Flint Basin

2. 11AA01

3. 13L180

4. 12M017

5. 08K001

6. 11K003

7. 12K014

8. 13J004

9. 08G001

10. 10G313

11.09F520

16. 11J011

Oconee Basin

12.21T001

Tennessee Basin

13.03PP01

Suwanee Basin

14. 19E009 17. 27E004

Ogeechee Basin

15.35P094




Figure 2. Area of use of principal aquifers and physiographic provinces in Georgia (modified from U.S. Geological Survey, 2006).


Groundwater Level Graphs


- For each of the 17 groundwater wells, EPD has prepared a graph that shows monthly average groundwater levels from January 2022 through December 2022;
- To help put these levels into perspective, for comparison purposes, each graph also shows:
 - Monthly average levels at that same well for the years 2007 and 2011 when groundwater levels were at or near recorded low levels across much of the state; and
 - And a statistical composite of historical conditions at that same gage showing the "lowest" 50, 20, 10, and 5 percent of all recorded monthly average levels at the same well.


How to Read the Groundwater Level Graphs Example: Well #11, 09F520, Flint River Basin


The groundwater level graph for Well #11, USGS 09F520 shows:


- The average monthly groundwater level in December 2022 was 49 ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in December have historically been lower than December 2022 about 25% of the time; about 75% of the time in December they have been higher.
- The average monthly groundwater level in December 2011 was 50.9 ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in December have historically been lower than December 2011 about 1% of the time; about 99% of the time in December they have been higher.
- The average monthly groundwater level in December 2007 was 50.9 ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in December have historically been lower than December 2007 about 1% of the time; about 99% of the time in December they have been higher.

































Back to Interpretation









Well #15, 35P094, Surficial Aquifer in Ogeechee Basin, Monthly Average Depth Below Land Surface







Reservoir Levels

Data Source: US Army Corps of Engineers

Coosa Basin

- 1. Carters
- 2. Allatoona

Chattahoochee Basin

- 3. Lanier
- 4. West Point
- 5. W.F. George

Savannah Basin


- 6. Hartwell
- 7. Thurmond





EPD monitors the water levels of seven reservoirs to assess drought conditions.


Reservoir Elevation Graphs


- The following graphs show the reservoir elevation curves for January 2022 through December 2022.
- Each graph also shows the Action Zone Divides (or Levels) for each reservoir
 - Zone 1 is the top layer of the conservation pool
 - Zone 2 is the layer below Zone 1
 - Zone 4 is the lowest layer in the conservation pool
 - There is no conservation storage below the bottom of Zone 4
- To put 2022 reservoir elevations into perspective, elevations for 2007 and 2011 are also shown.












Back to Map


84




2023 ACF Basin Composite Conservation and Flood Storage

Actual data thru 1/9/2023

Add value of 1,856,000 acre-ft to include inactive storage.





Climate Prediction Center 3-month Temperature and Precipitation Probability Outlook and Seasonal Drought Outlook

> Data Source: http://www.cpc.ncep.noaa.gov/

