# **Drought Indicators Report**

### Georgia Environmental Protection Division March 2017

## Background

Pursuant to the Rules for Drought Management, <u>Section 391-3-3-.04 Drought</u> <u>Indicators and Triggers</u>, the Director of EPD monitors climatic indicators and water supply conditions to assess drought occurrence and severity, and its impact upon the ability of public water systems to provide adequate supplies of water. These indicators and conditions may include, but not be limited, to the following:

- U.S. Drought Monitor;
- Precipitation;
- Streamflow;
- Groundwater;
- Reservoir levels;
- Short term climate predictions;
- Soil moisture; and
- Water supply conditions.

## Background

- The Rules require EPD to report on current climatic indicators at least semi-annually or monthly when any part of the state has experienced at least two consecutive months of severe drought.
- This reports compare current conditions to historical levels (and/or reservoir rule curves) for each of the following indicators:
  - Precipitation during the prior 3, 6, and 12 months;
  - Streamflow at the select United States Geological Survey gages;
  - Groundwater levels at select United States Geological Survey monitoring wells; and
  - Reservoir levels at Allatoona Lake, Lake Hartwell, Clarks Hill Lake, and Lake Lanier.
- The following sections of this presentation provide the data and information sources analyzed by EPD in developing this drought indicators report for conditions as of March 20, 2017.

### Drought Indicator Analysis Summary (slide 1 of 2)

<u>U.S. Drought</u> Monitor - Extreme drought is in all or parts of 17 counties in the northern portion of the state. Severe drought remains in an additional 33 counties north of the fall line. This week marks the 42nd week of continuous severe (or more intense) drought in northwest Georgia, the 40th week for the northern Atlanta metro area, the 39th week in parts of the northeast.

**Precipitation -** The 3 month records show surplus amounts of rainfall along and around the fall line and southwest of metro Atlanta, while deficits generally cover the remainder of the state. North central and northeast Georgia still show the most deficits, with up to -12" in the worst areas. The 6 month records show areas north of the fall line with the worst deficits in the state, with extreme north central and northeast Georgia showing deficits greater than -20" in the worst areas. Slight deficits are emerging in central Georgia and extreme south Georgia also in the 6 month time period. The 12 month records show near normal to a slight surplus of rainfall in southwest and parts of southeast Georgia, while the greatest deficits still exist north of the fall line. Central and parts of southeast Georgia still show slight deficits in the long term.

<u>Soil Moisture</u> - The vast majority of state shows normal to some level of dryness, with the worst conditions in much of northeast Georgia with exceptional dryness, and in south central to southeast Georgia where up to moderate dryness is present. North Georgia still remains in severe drought in the long term periods due to the longevity of the drought in those core drought areas. Dry conditions in north Florida continue to creep northward into south Georgia. West-central areas of the state generally show normal soil moisture conditions.

### Drought Indicator Analysis Summary (slide 2 of 2)

**Stream Flows** - 25 of the 34 observation sites are at or below 2007 and/or 2011 levels and many locations show January levels well above normal. 19 gages show flows at or lower than the 5th percentile.

<u>**Groundwater</u>** - Groundwater levels vary by location. 5 of the 14 of the monitoring wells EPD uses to track drought conditions are below median levels. 12 are above the 20th percentile and 2are below the 5th percentile of the historical record.</u>


**<u>Reservoir Levels</u>** - In the ACT, Allatoona is at its current rule curve, which is rising toward its summer normal pool level and Carters is less than 3.5 feet from its rule curve and within Zone 1. In the ACF, Lanier is zone 4, WestPoint and George both remain above their respective rule curves. ACF Composite storage is in Zone 3. In the Savannah Basin, both Hartwell and Thurmond are half way between elevation Levels 2 and 3and remain in Corps drought level 2 operations.

Short Term Climate Prediction - Drought conditions in north Georgia are predicted to persist.

<u>Water Supplies</u> - Many systems are reporting that local water supplies have recovered or nearly recovered. Lanier is the primary exception, and it is approximately 10 feet down. Systems are still generally advising a cautious approach to discretionary water use.

# **US Drought Monitor**


Data Source: http://droughtmonitor.unl.edu/



### U.S. Drought Monitor Georgia


#### March 14, 2017 (Released Thursday, Mar. 16, 2017) Valid 8 a.m. EDT

Drought Conditions (Percent Area)



	None	D0-D4	D1-D4	D2-D4	D3-D4	D4
Current	68.50	31.50	26.59	19.08	6.94	0.00
Last Week 3/7/2017	66.26	33.74	27.56	19.39	8.69	0.00
3 Month s Ago 12/13/2016	9.09	90.91	83.09	63.26	50.18	27.25
Start of Calend ar Year 1/3/2017	11.31	88.69	73.48	39.33	19.28	0.00
Start of Water Year 9/27/2016	35.37	64.63	45.84	34.50	14.67	1.58
One Year Ago 3/15/2016	86.78	13.22	0.00	0.00	0.00	0.00

#### Intensity:



D3 Extreme Drought

D4 Exceptional Drought

D2 Severe Drought

D1 Moderate Drought

The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. See accompanying text summary for forecast statements.

#### Author:

Brian Fuchs National Drought Mitigation Center




#### http://droughtmonitor.unl.edu/


# 3, 6, and 12 Month Percent of Normal Precipitation

Data Source: http://climate.ncsu.edu/drought


## 3 Month Percent of Normal Precipitation



## 6 Month Percent of Normal Precipitation




## 12 Month Percent of Normal Precipitation



# Soil Moisture Conditions

Data Source: http://www.hydro.washington.edu/forecast/monitor/curr/con us.mexico/east.vic.sm\_qnt.gif



VIC Soil Moisture Percentiles (wrt/ 1916-2004) Eastern United States - 20170318


Current (daily updated) percentiles for soil moisture (SWE) with respect to the climatological period (1916-2004).

# **Streamflow Conditions**

Data Source: USGS

## **Streamflow Monitoring**

- As shown on the following slide, EPD Monitors 34 USGS stream gages in 13 of the State's major river basins to assess drought conditions.
- These gages were selected because each has:
  - Long-term and relatively complete records for recent decades; and
  - Relatively low consumptive water use implications and streamflows are not heavily influenced by dams.
- Note: Hydrologic conditions of major rivers with streamflows that are heavily influenced by dams can be assessed by reviewing status of major storage reservoirs



#### USGS Stream Gages Monitored by EPD to Assess Drought Conditions

GAGE#	BASIN	GAGE NAME
1	TENNESSEE	LOOKOUT CREEK NEAR NEW ENGLAND
2	TENNESSEE	NOTTELY RIVER NEAR BLAIRSVILLE
3	COOSA	CHATTOOGA RIVER AT SUMMERVILLE
4	COOSA	TALKING ROCK CREEK NEAR HINTON
5	COOSA	ETOWAH RIVER AT CANTON
6	CHATTAHOOCHEE	CHATTAHOOCHEE RIVER AT CORNELIA
7	CHATTAHOOCHEE	CHESTATEE RIVER NEAR DAHLONEGA
8	CHATTAHOOCHEE	NEW RIVER AT GA 100 NEAR CORINTH
9	CHATTAHOOCHEE	UPATOI CREEK AT COLUMBUS
10	FLINT	FLINT RIVER AT GA26 NEAR MONTEZUMA
11	FLINT	FLINT RIVER AT ALBANY
12	FLINT	ICHAWAYNOCHAWAY CREEK AT MILFORD
13	FLINT	SPRING CREEK NEAR IRON CITY
14	OCMULGEE	ALCOVY RIVER ABOVE COVINGTON
15	OCMULGEE	OCMULGEE RIVER AT MACON
16	OCMULGEE	TOBESOFKEE CREEK NEAR MACON
17	OCMULGEE	TUCSAWHATCHEE CREEK NEAR
		HAWKINSVILLE
18	OCONEE	MIDDLE OCONEE RIVER NEAR ATHENS
19	OCONEE	LITTLE RIVER NEAR EATONTON
20	OCONEE	OCONEE RIVER AT DUBLIN
21	ALTAMAHA	OHOOPEE RIVER NEAR REIDSVILLE
22	SAVANNAH	CHATTOOGA RIVER NEAR CLAYTON
23	SAVANNAH	BROAD RIVER NEAR BELL
24	SAVANNAH	BEAVERDAM CREEK NEAR SARDIS
25	SAVANNAH	BRIER CREEK AT MILLHAVEN
26	OGEECHEE	CANOOCHEE RIVER NEAR CLAXTON
27	OGEECHEE	OGEECHEE RIVER NEAR EDEN
28	OCHLOCKONEE	OCHLOCKONEE RIVER NEAR THOMASVILLE
29	SUWANEE	WITHLACOOCHEE RIVER NEAR PINETTA FL
30	SUWANEE	ALAPAHA RIVER AT STATENVILLE
31	SUWANEE	SUWANNEE RIVER AT US 441, AT FARGO
32	SATILLA	SATILLA RIVER NEAR WAYCROSS
33	SATILLA	LITTLE SATILLA RIVER NEAR OFFERMAN
34	ST MARY	ST MARYS RIVER NEAR MACCLENNY FL

## Streamflow Graphs

- For each of the 34 gages, EPD has prepared a graph that shows monthly average streamflow from January, 2017 through February, 2017;
- To help put these streamflow conditions into perspective, for comparison purposes, each graph also shows:
  - Monthly average streamflows for the years 2007 and 2011 when streamflows were at or near recorded low levels across much of the state; and
  - A statistical composite of historical conditions showing the "driest" 50, 20, 10, and 5 percent of all recorded monthly average stream flows at the same gage.

## How to Read the Streamflow Graphs Example #1: Etowah River at Canton


The streamflow graph for Gage #5, <u>USGS Etowah River gage at Canton</u> shows :


- Average stream flow for February 2017 was 428 cfs. The statistical composite of all historical data for this gage shows that average streamflow in February has historically been lower than February 2017 about 0% of the time; about 100% of the time in February it has been higher.
- Average stream flow in February 2011 was 932 cfs. The statistical composite of all historical data for this gage shows that average streamflow for February has historically been lower than February 2011 only 15% of the time; 85% of the time in February it has been higher.
- Average stream flow in February 2007 was 885 cfs. The statistical composite of all historical data for this gage shows that average streamflow for February has historically been lower than February 2007 only 15% of the time; 85% of the time in February it has been higher.


## How to Read the Streamflow Graphs <u>Example #2:</u> Flint River at Albany


The streamflow graph for Gage #11, <u>USGS Flint River gage at Albany</u> shows:


- Average stream flow for February 2017 was 5,792 cfs. The statistical composite of all historical data for this gage shows that average streamflow in February has historically been lower than February 2017 about 20% of the time; about 80% of the time in February it has been higher.
- Average stream flow in February 2011 was 6,534 cfs. The statistical composite of all historical data for this gage shows that average streamflow for February has historically been lower than February 2011 about 25% of the time; about 75% of the time in February it has been higher.
- Average stream flow in February 2007 was 5,591 cfs. The statistical composite of all historical data for this gage shows that average streamflow for February has historically been lower than February 2007 about 15% of the time; about 85% of the time in February it has been higher.


































































































# **Groundwater Levels**

Data Source: USGS

### Rationale for Choosing USGS Monitoring Wells

EPD monitors 14 groundwater USGS monitoring wells shown on the following slide to assess drought conditions. These wells were selected for monitoring because they have:

- Long-term monitoring records consisting of three decades or more of data; and
- Real-time monitoring that represents the most up-to-date conditions.

### USGS Wells Monitored by EPD to Assess Drought Conditions

#### Savannah Basin

1.30AA04

#### **Flint Basin**

2. 11AA01

- 3. 13L180
- 4. 12M017
- 5. 08K001
- 6. 11K003
- 7. 12K014
- 8. 13J004
- 9. 08G001
- 10. 10G313
- 11.09F520

#### **Oconee Basin**

12. 21T001

#### Altamaha Basin

13. 26R001

#### Suwanee Basin

14. 19E009




Figure 2. Area of use of principal aquifers and physiographic provinces in Georgia (modified from U.S. Geological Survey, 2006).


### Groundwater Level Graphs


- For each of the 15 groundwater wells, EPD has prepared a graph that shows monthly average groundwater levels from January, 2017 through February, 2017;
- To help put these levels into perspective, for comparison purposes, each graph also shows:
  - Monthly average levels at that same well for the years 2007 and 2011 when groundwater levels were at or near recorded low levels across much of the state; and
  - And a statistical composite of historical conditions at that same gage showing the "lowest" 50, 20, 10, and 5 percent of all recorded monthly average levels at the same well.


## How to Read the Groundwater Level Graphs Example: Well #11, 09F520, Flint River Basin


The groundwater level graph for Well #11, USGS 09F520 shows:

- The average monthly groundwater level for February 2017 was 46ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in February have historically been lower than February 2017 about 50% of the time; about 50% of the time in February they have been higher.
- The average monthly groundwater level in February 2011 was 49ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in February have historically been lower than February 2017 about 15% of the time; about 85% of the time in February they have been higher.
- The average monthly groundwater level in February 2007 was 46.8ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in February have historically been lower than February 2017 about 40% of the time; about 60% of the time in February they have been higher.













Back to Map








Back to Map




















# **Reservoir Levels**

Data Source: US Army Corps of Engineers

### Coosa Basin

- 1. Carters
- 2. Allatoona

### **Chattahoochee Basin**

- 3. Lanier
- 4. West Point
- 5. W.F. George

### Savannah Basin


- 6. Hartwell
- 7. Thurmond





EPD monitors the water levels of seven reservoirs to assess drought conditions.


# **Reservoir Elevation Graphs**


- The following graphs show the reservoir elevation curves for January, 2017 through February, 2017.
- Each graph also shows the Action Zone Divides (or Levels) for each reservoir
  - Zone 1 is the top layer of the conservation pool
  - Zone 2 is the layer below Zone 1
  - Zone 4 is the lowest layer in the conservation pool
  - There is no conservation storage below the bottom of Zone 4
- To put 2017 reservoir elevations into perspective, elevations for 2007 and 2011 are also shown.
















#### 2017 ACF Basin Composite Conservation and Flood Storage

Actual data thru 3-15-2017

Add value of 1,856,000 acre-ft to include inactive storage.


Compiled by USACOE.






Climate Prediction Center 3-month Temperature and Precipitation Probability Outlook and Seasonal Drought Outlook

> Data Source: http://www.cpc.ncep.noaa.gov/




## **Precipitation Outlook**



### **U.S. Seasonal Drought Outlook** Drought Tendency During the Valid Period

### Valid for March 16 - June 30, 2017 Released March 16, 2017



Depicts large-scale trends based on subjectively derived probabilities guided by short- and long-range statistical and dynamical forecasts. Use caution for applications that can be affected by short lived events. "Ongoing" drought areas are based on the U.S. Drought Monitor areas (intensities of D1 to D4).

NOTE: The tan areas imply at least a 1-category improvement in the Drought Monitor intensity levels by the end of the period, although drought will remain. The green areas imply drought removal by the end of the period (D0 or none).

#### **Drought persists**

**Drought remains but improves** 

**Drought removal likely** 

**Drought development likely** 



### http://go.usa.gov/3eZ73