

# Drought Indicators Report

Georgia Environmental Protection Division  
May 2017

# Background

Pursuant to the Rules for Drought Management, Section 391-3-3-.04 Drought Indicators and Triggers, the Director of EPD monitors climatic indicators and water supply conditions to assess drought occurrence and severity, and its impact upon the ability of public water systems to provide adequate supplies of water. These indicators and conditions may include, but not be limited, to the following:

- U.S. Drought Monitor;
- Precipitation;
- Streamflow;
- Groundwater;
- Reservoir levels;
- Short term climate predictions;
- Soil moisture; and
- Water supply conditions.

# Background

- The Rules require EPD to report on current climatic indicators at least semi-annually or monthly when any part of the state has experienced at least two consecutive months of severe drought.
- This reports compare current conditions to historical levels (and/or reservoir rule curves) for each of the following indicators:
  - Precipitation during the prior 3, 6, and 12 months;
  - Streamflow at the select United States Geological Survey gages;
  - Groundwater levels at select United States Geological Survey monitoring wells; and
  - Reservoir levels at Allatoona Lake, Lake Hartwell, Clarks Hill Lake, and Lake Lanier.
- The following sections of this presentation provide the data and information sources analyzed by EPD in developing this drought indicators report for conditions as of May 12, 2017.

# Drought Indicator Analysis Summary (slide 1 of 2)

- **U.S. Drought Monitor** – Extreme drought is in parts of 3 counties in the northern portion of the state. Severe drought remains in parts of an additional 7 counties north of the fall line. Drought conditions have expanded in south Georgia, with extreme drought conditions in parts of 12 counties and severe drought conditions in 32 counties.
- **Precipitation**– The 3 month records show the majority of the state with below normal precipitation except areas of northwest and northeast Georgia, where surplus amounts of rainfall are present. Conditions in much of north Georgia are near to slightly below normal, with worsening conditions in the southern half of the state. Areas of south Georgia along and north of the Florida border show the worst deficits of up to -12”, while areas of central Georgia are gradually worsening. The 6 month records show a surplus of rainfall in extreme northwest, southwest, east central Georgia, and just south of the metro Atlanta area, while extreme northeast and southeast Georgia show deficits up to 12”. The rest of the state shows normal rainfall in the 6 month time period. The 12 month records show much of the state with below normal precipitation, particularly in the northern half of the state and along the Florida border, while areas of southeast Georgia show near to slightly above normal precipitation. Northeast Georgia shows the worst deficits in the longer term 12-month time period with around 50% of normal precipitation.
- **Soil Moisture**– Dry to extremely dry soil moisture conditions exist in the southern half of the state south of metro Atlanta except for west central Georgia and along areas along the northern half of the coast where normal conditions are present. Conditions northward show normal to exception wetness, particularly in northwest and northeast Georgia. Areas in southeast Georgia show the most exceptional short and longer term dryness, with central and southwest Georgia close behind.

## Drought Indicator Analysis Summary (slide 2 of 2)


- **Stream Flows** - 29 of the 34 observation sites are at or below 2007 and/or 2011 level. 2 gages show flows at or lower than the 5th percentile.
- **Groundwater** - Groundwater levels vary by location. 12 of the 14 of the monitoring wells EPD uses to track drought conditions are below median levels. 12 are at or above the 20th percentile and 1 is below the 5th percentile of the historical record. Data for one of the 14 wells was unavailable at the time of this report.
- **Reservoir Levels** - In the ACT, Allatoona and Carters are both at or above rule curve. In the ACF, Lanier is zone 4, WestPoint is in zone 1 and George remains above its rule curve. ACF Composite storage is in Zone 2. The ACF remains in Corps drought operations. In the Savannah Basin, both Hartwell and Thurmond are both in Level 3 and remain in Corps drought level 2 operations.
- **Short Term Climate Prediction** - Drought conditions in North and South Georgia are predicted to alleviate, conditions in central Georgia are predicted to persist.
- **Water Supplies** - Many systems are reporting that local water supplies have recovered or nearly recovered. Lanier is the primary exception, and it is approximately 8 feet down. Systems are still generally advising a cautious approach to discretionary water use.

# US Drought Monitor

Data Source:  
<http://droughtmonitor.unl.edu/>

# U.S. Drought Monitor

May 16, 2017  
(Released Thursday, May. 18, 2017)  
Valid 8 a.m. EDT





*Author:*  
Brad Rippey  
U.S. Department of Agriculture

***Drought Impact Types:***  
~ Delineates dominant impacts  
S = Short-Term, typically less than 6 months (e.g. agriculture, grasslands)  
L = Long-Term, typically greater than 6 months (e.g. hydrology, ecology)

***Intensity:***  
D0 Abnormally Dry  
D1 Moderate Drought  
D2 Severe Drought  
D3 Extreme Drought  
D4 Exceptional Drought

The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. See accompanying text summary for forecast statements.



<http://droughtmonitor.unl.edu/>

# U.S. Drought Monitor

## Georgia


May 16, 2017

(Released Thursday, May. 18, 2017)

Valid 8 a.m. EDT

*Drought Conditions (Percent Area)*

	None	D0-D4	D1-D4	D2-D4	D3-D4	D4
<b>Current</b>	11.83	88.17	54.01	23.19	7.21	0.00
<b>Last Week</b> <i>05-09-2017</i>	11.87	88.13	54.20	23.47	6.24	0.00
<b>3 Months Ago</b> <i>02-14-2017</i>	63.07	36.93	30.94	19.79	5.39	0.00
<b>Start of Calendar Year</b> <i>01-03-2017</i>	11.31	88.69	73.48	39.33	19.28	0.00
<b>Start of Water Year</b> <i>09-27-2016</i>	35.37	64.63	45.84	34.50	14.67	1.58
<b>One Year Ago</b> <i>05-17-2016</i>	52.92	47.08	27.04	0.00	0.00	0.00



Intensity:

- D0 Abnormally Dry
- D1 Moderate Drought
- D2 Severe Drought
- D3 Extreme Drought
- D4 Exceptional Drought

*The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. See accompanying text summary for forecast statements.*

Author:

Brad Rippey  
U.S. Department of Agriculture






# 3, 6, and 12 Month Percent of Normal Precipitation


Data Source:

<http://climate.ncsu.edu/drought>

# 3 Month Percent of Normal Precipitation




90 day SPI for May 11, 2017




Transparency:   
[About these maps](#)


# 6 Month Percent of Normal Precipitation




6 month SPI for May 11, 2017



# 12 Month Percent of Normal Precipitation



12 month SPI for May 11, 2017



Transparency:




[About these maps](#)

# Soil Moisture Conditions

Data Source:

[http://www.hydro.washington.edu/forecast/monitor/curr/conus.mexico/east.vic.sm\\_qnt.gif](http://www.hydro.washington.edu/forecast/monitor/curr/conus.mexico/east.vic.sm_qnt.gif)

VIC Soil Moisture Percentiles (wrt' 1916-2004)  
Eastern United States - 20170511



Current (daily updated) percentiles for soil moisture (SWE) with respect to the climatological period (1916-2004).

# Streamflow Conditions

Data Source: USGS


# Streamflow Monitoring

- As shown on the following slide, EPD Monitors 34 USGS stream gages in 13 of the State's major river basins to assess drought conditions.
- These gages were selected because each has:
  - Long-term and relatively complete records for recent decades; and
  - Relatively low consumptive water use implications and streamflows are not heavily influenced by dams.
- Note: Hydrologic conditions of major rivers with streamflows that are heavily influenced by dams can be assessed by reviewing status of major storage reservoirs



# Georgia's 14 River Basins

## USGS Stream Gages Monitored by EPD to Assess Drought Conditions



<u>GAGE#</u>	<u>Basin</u>	<u>GAGE NAME</u>
1	TENNESSEE	LOOKOUT CREEK NEAR NEW ENGLAND
2	TENNESSEE	NOTTELY RIVER NEAR BLAIRSVILLE
3	COOSA	CHATTOOGA RIVER AT SUMMERVILLE
4	COOSA	TALKING ROCK CREEK NEAR HINTON
5	COOSA	ETOWAH RIVER AT CANTON
6	CHATTAHOOCHEE	CHATTAHOOCHEE RIVER AT CORNELIA
7	CHATTAHOOCHEE	CHESTATEE RIVER NEAR DAHLONEGA
8	CHATTAHOOCHEE	NEW RIVER AT GA 100 NEAR CORINTH
9	CHATTAHOOCHEE	UPATOI CREEK AT COLUMBUS
10	FLINT	FLINT RIVER AT GA26 NEAR MONTEZUMA
11	FLINT	FLINT RIVER AT ALBANY
12	FLINT	ICHAWAYNOCHAWAY CREEK AT MILFORD
13	FLINT	SPRING CREEK NEAR IRON CITY
14	OCMULGEE	ALCOVY RIVER ABOVE COVINGTON
15	OCMULGEE	OCMULGEE RIVER AT MACON
16	OCMULGEE	TOBESOFKEE CREEK NEAR MACON
17	OCMULGEE	TUCSAWHATCHEE CREEK NEAR HAWKINSVILLE
18	OCONEE	MIDDLE OCONEE RIVER NEAR ATHENS
19	OCONEE	LITTLE RIVER NEAR EATONTON
20	OCONEE	OCONEE RIVER AT DUBLIN
21	ALTAMAHA	OHOOPPEE RIVER NEAR REIDSVILLE
22	SAVANNAH	CHATTOOGA RIVER NEAR CLAYTON
23	SAVANNAH	BROAD RIVER NEAR BELL
24	SAVANNAH	BEAVERDAM CREEK NEAR SARDIS
25	SAVANNAH	BRIER CREEK AT MILLHAVEN
26	OGEECHEE	CANOOCHEE RIVER NEAR CLAXTON
27	OGEECHEE	OGEECHEE RIVER NEAR EDEN
28	OCHLOCKONEE	OCHLOCKONEE RIVER NEAR THOMASVILLE
29	SUWANEЕ	WITHLACOOCHEE RIVER NEAR PINETTA FL
30	SUWANEЕ	ALAPAHA RIVER AT STATENVILLE
31	SUWANEЕ	SUWANNEE RIVER AT US 441, AT FARGO
32	SATILLA	SATILLA RIVER NEAR WAYCROSS
33	SATILLA	LITTLE SATILLA RIVER NEAR OFFERMAN
34	ST MARY'S	ST MARYS RIVER NEAR MACCLENNY FL

# Streamflow Graphs

- For each of the 34 gages, EPD has prepared a graph that shows monthly average streamflow from January, 2017 through April, 2017;
- To help put these streamflow conditions into perspective, for comparison purposes, each graph also shows:
  - Monthly average streamflows for the years 2007 and 2011 when streamflows were at or near recorded low levels across much of the state; and
  - A statistical composite of historical conditions showing the “driest” 50, 20, 10, and 5 percent of all recorded monthly average stream flows at the same gage.

# How to Read the Streamflow Graphs

## Example #1: [Etowah River at Canton](#)

The streamflow graph for Gage #5, [USGS Etowah River gage at Canton](#) shows :

- Average stream flow for April 2017 was 849 cfs. The statistical composite of all historical data for this gage shows that average streamflow in April has historically been lower than April 2017 about 10% of the time; about 90% of the time in April it has been higher.
- Average stream flow in April 2011 was 1604 cfs. The statistical composite of all historical data for this gage shows that average streamflow for April has historically been lower than April 2011 only 50% of the time; 50% of the time in April it has been higher.
- Average stream flow in April 2007 was 746 cfs. The statistical composite of all historical data for this gage shows that average streamflow for April has historically been lower than April 2007 only 5% of the time; 95% of the time in April it has been higher.


# How to Read the Streamflow Graphs

## Example #2: Flint River at Albany

The streamflow graph for Gage #11, [USGS Flint River gage at Albany](#) shows:


- Average stream flow for April 2017 was 4,734 cfs. The statistical composite of all historical data for this gage shows that average streamflow in April has historically been lower than April 2017 about 20% of the time; about 80% of the time in April it has been higher.
- Average stream flow in April 2011 was 4,789 cfs. The statistical composite of all historical data for this gage shows that average streamflow for April has historically been lower than April 2011 about 20% of the time; about 80% of the time in April it has been higher.
- Average stream flow in April 2007 was 3,310 cfs. The statistical composite of all historical data for this gage shows that average streamflow for April has historically been lower than April 2007 about 5% of the time; about 95% of the time in April it has been higher.

## Gage #1. USGS #03568933, Tennessee Basin, LOOKOUT CREEK NEAR NEW ENGLAND, GA




[Back to Map](#)

## Gage #2, USGS #03550500, Tennessee Basin, NOTTELY RIVER NEAR BLAIRSVILLE, GA




[Back to Map](#)

### Gage #3. USGS #02398000, Coosa Basin, Chattooga River at Summerville, GA



[Back to Map](#)


# Gage #4, USGS #02382200, Coosa Basin, TALKING ROCK CREEK NEAR HINTON, GA



[Back to Map](#)




# Gage #5, USGS #02392000, Coosa Basin, Etowah River at Canton, GA



[Back to Interpretation](#)


[Back to Map](#)

# Gage #6, USGS #02331600, Chatthoochee Basin, CHATTAHOOCHEE RIVER AT CORNELIA, GA




[Back to Map](#)

## Gage #7, USGS #02333500, Chatahoochee Basin, CHESTATEE RIVER NEAR DAHLONEGA, GA




[Back to Map](#)

**Gage #8, USGS #02338660, Chattahoochee Basin,  
NEW RIVER AT GA 100, NEAR CORINTH**




[Back to Map](#)

## Gage #9, USGS #02341800, Chattahoochee Basin, UPATOI CREEK NEAR COLUMBUS, GA




[Back to Map](#)

## Gage #10. USGS #02349605, Flint Basin, FLINT RIVER AT GA26 NEAR MONTEZUMA, GA



[Back to Map](#)


## Gage #11, USGS #02352500, Flint Basin, FLINT RIVER AT ALBANY, GA



[Back to Interpretation](#)

[Back to Map](#)


## Gage #12. USGS #02353500, Flint Basin, Ichawaynochaway Creek at Milford, GA



[Back to Map](#)




# Gage #13. USGS #02357000, Flint River, Spring Creek near Iron City, GA




[Back to Map](#)

## Gage #14. USGS #02208450, Ocumulgee Basin, ALCOVY RIVER above COVINGTON, GA




[Back to Map](#)

## Gage #15. USGS #02213000, Ocmulgee Basin, Ocmulgee River at Macon, GA




[Back to Map](#)

## Gage #16. USGS #02213500, Ocmulgee Basin, TOBESOFKEE CREEK near MACON, GA




[Back to Map](#)

## Gage #17. USGS #02215100, Ocmulgee Basin, TUCSAWHATCHEE CREEK near HAWKINSVILLE, GA




[Back to Map](#)

# Gage #18. USGS #02217500, Oconee Basin, MIDDLE OCONEE RIVER near ATHENS, GA




# Gage #19. USGS #02220900, Oconee Basin, LITTLE RIVER near EATONTON, GA



[Back to Map](#)


## Gage #20. USGS #02223500, Oconee Basin, Oconee River at Dublin, GA



[Back to Map](#)




# Gage #21. USGS #02225500, Altamaha Basin, OHOOPEE RIVER near REIDSVILLE, GA




[Back to Map](#)

## Gage #22. USGS #02177000, Savannah Basin, CHATTOOGA RIVER near CLAYTON, GA




[Back to Map](#)

# Gage #23. USGS #02192000, Savannah Basin Broad River near Bell, GA




[Back to Map](#)

## Gage #24. USGS #02198100, Savannah Basin, BEAVERDAM CREEK near SARDIS, GA




[Back to Map](#)

## Gage #25. USGS #02198000 , Savannah Basin, BRIER CREEK at MILLHAVEN, GA




[Back to Map](#)

## Gage #26. USGS #02203000, Ogeechee Basin, CANOOCHEE RIVER near CLAXTON, GA




[Back to Map](#)

## Gage #27. USGS #02202500, Ogeechee Basin, Ogeechee River near Eden, GA




[Back to Map](#)

## Gage #28. USGS #02327500, Ochlockonee Basin, OCHLOCKONEE RIVER near THOMASVILLE, GA






## Gage #29. USGS #02319000, Suwanee Basin, WITHLACOOCHEE RIVER near PINETTA, FL




[Back to Map](#)

# Gage #30. USGS #02317500, Suwanee Basin, Alapaha River at Statenville, GA




[Back to Map](#)

## Gage #31. USGS #02314500, Suwannee Basin, SUWANNEE RIVER AT US 441, AT FARGO, GA




[Back to Map](#)

# Gage #32. USGS #02226500, Satilla Basin, Satilla River near Waycross, GA




[Back to Map](#)

## Gage #33. USGS #02227500, Satilla Basin, LITTLE SATILLA RIVER near OFFERMAN, GA



[Back to Map](#)

## Gage #34. USGS #02231000, St Mary Basin, ST. MARYS RIVER near MACCLENNY, FL



[Back to Map](#)

# Groundwater Levels

Data Source: USGS

## Rationale for Choosing USGS Monitoring Wells

EPD monitors 14 groundwater USGS monitoring wells shown on the following slide to assess drought conditions. These wells were selected for monitoring because they have:

- Long-term monitoring records consisting of three decades or more of data; and
- Real-time monitoring that represents the most up-to-date conditions.



# USGS Wells Monitored by EPD to Assess Drought Conditions

## Savannah Basin

1. 30AA04

## Flint Basin

- 2. 11AA01
- 3. 13L180
- 4. 12M017
- 5. 08K001
- 6. 11K003
- 7. 12K014
- 8. 13J004
- 9. 08G001
- 10. 10G313
- 11. 09F520

## Oconee Basin


12. 21T001

## Altamaha Basin

13. 26R001

## Suwanee Basin

14. 19E009



### EXPLANATION

<b>Coastal Plain aquifers</b>	<b>Piedmont and Blue Ridge aquifers</b>
Surficial aquifer system—Not a principal aquifer. Shown on block only	Crystalline-rock aquifers
Floridan aquifer system	<b>Valley and Ridge and Appalachian Plateau aquifers</b>
Claiborne and Clayton aquifers	Paleozoic-rock aquifers
Cretaceous aquifer system	

Figure 2. Area of use of principal aquifers and physiographic provinces in Georgia (modified from U.S. Geological Survey, 2006).

# Groundwater Level Graphs

- For each of the 15 groundwater wells, EPD has prepared a graph that shows monthly average groundwater levels from January, 2017 through April, 2017;
- To help put these levels into perspective, for comparison purposes, each graph also shows:
  - Monthly average levels at that same well for the years 2007 and 2011 when groundwater levels were at or near recorded low levels across much of the state; and
  - And a statistical composite of historical conditions at that same gage showing the “lowest” 50, 20, 10, and 5 percent of all recorded monthly average levels at the same well.


# How to Read the Groundwater Level Graphs

## Example: [Well #11, 09F520, Flint River Basin](#)

The groundwater level graph for Well #11, USGS 09F520 shows:


- The average monthly groundwater level for April 2017 was 48ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in April have historically been lower than April 2017 about 10% of the time; about 90% of the time in April they have been higher.
- The average monthly groundwater level in April 2011 was 49ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in April have historically been lower than April 2017 about 2% of the time; about 98% of the time in April they have been higher.
- The average monthly groundwater level in April 2007 was 47.4ft below land surface. The statistical composite of all historical data for this well shows that monthly average groundwater levels in April have historically been lower than April 2017 about 10% of the time; about 90% of the time in April they have been higher.

# Well #1, 30AA04, Gordon & Dublin Aquifers in Savannah Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

# Well #2, 11AA01, Surficial Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

## Well #3, 13L180, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

## Well #4, 12M017, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface



[Back to Map](#)


## Well #5, 08K001, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface



[Back to Map](#)




## Well #6, 11K003, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

## Well #7, 12K014, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

## Well #8, 13J004, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

# Well #9, 08G001, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface




## Well #10, 10G313, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface



[Back to Map](#)


## Well #11, 09F520, Floridan Aquifer in Flint Basin, Monthly Average Depth Below Land Surface



[Back to Interpretation](#)


[Back to Map](#)

## Well #12, 21T001, Floridan Aquifer in Oconee Basin, Monthly Average Depth Below Land Surface



[Back to Map](#)

## Well #13, 26R001, Floridan Aquifer in Altamaha Basin, Monthly Average Depth Below Land Surface




[Back to Map](#)

Data unavailable for this time period



# Well #14, 19E009, Floridan Aquifer in Suwanee Basin, Monthly Average Depth Below Land Surface



[Back to Map](#)

# Reservoir Levels

Data Source:  
US Army Corps of Engineers

### Coosa Basin

1. Carters
2. Allatoona

### Chattahoochee Basin

3. Lanier
4. West Point
5. W.F. George

### Savannah Basin

6. Hartwell
7. Thurmond




EPD monitors the water levels of seven reservoirs to assess drought conditions.

# Reservoir Elevation Graphs


- The following graphs show the reservoir elevation curves for January, 2017 through April, 2017.
- Each graph also shows the Action Zone Divides (or Levels) for each reservoir
  - Zone 1 is the top layer of the conservation pool
  - Zone 2 is the layer below Zone 1
  - Zone 4 is the lowest layer in the conservation pool
  - There is no conservation storage below the bottom of Zone 4
- To put 2017 reservoir elevations into perspective, elevations for 2007 and 2011 are also shown.

# CARTERS ELEVATION




[Back to Map](#)

# ALLATOONA ELEVATION




# LAKE LANIER ELEVATION



[Back to Map](#)


# WEST POINT ELEVATION



[Back to Map](#)




# W.F.GEORGE ELEVATION



[Back to Map](#)


## 2017 ACF Basin Composite Conservation and Flood Storage




Actual data thru 5-9-2017

Add value of 1,856,000 acre-ft to include inactive storage.

# Lake HARTWELL ELEVATION



# LAKE CLARK HILL (THURMOND) ELEVATION




# Climate Prediction Center 3-month Temperature and Precipitation Probability Outlook and Seasonal Drought Outlook


Data Source:

<http://www.cpc.ncep.noaa.gov/>

# Temperature Outlook




# Precipitation Outlook



# U.S. Seasonal Drought Outlook

## Drought Tendency During the Valid Period





Valid for May 18 - August 31, 2017  
Released May 18, 2017




Depicts large-scale trends based on subjectively derived probabilities guided by short- and long-range statistical and dynamical forecasts. Use caution for applications that can be affected by short lived events. "Ongoing" drought areas are based on the U.S. Drought Monitor areas (intensities of D1 to D4).

NOTE: The tan areas imply at least a 1-category improvement in the Drought Monitor intensity levels by the end of the period, although drought will remain. The green areas imply drought removal by the end of the period (D0 or none).

Author:  
R. Tinker  
NOAA/NWS/NCEP/Climate Prediction Center

-  Drought persists
-  Drought remains but improves
-  Drought removal likely
-  Drought development likely



<http://go.usa.gov/3eZ73>